Proceedings of ICBioMed 2024 Workshop: Computational Proteomics in Drug Discovery and Development from Medicinal Plants
DOI: 10.54254/2753-8818/75/2024.LA19353

The relationship between obesity and gut microbiota

Simiao Cai

Qingdao No.2 Middle School, Shandong Province, China

jungkook1015111@outlook.com

Abstract. This article begins by discussing the important the impact of the gut microbiome on obesity, emphasizing how changes in microbial diversity and composition can influence fat storage, energy extraction, and metabolic health. It first outlines the connection between gut microbes and key obesity-related mechanisms such as fat accumulation, insulin resistance, and systemic inflammation. Next, external elements like nutrition, medications, lifestyle, and genetics that influence the composition of the gut microbiome and, consequently, obesity outcomes are examined. Afterward, potential interventions for restoring microbial balance are discussed, including probiotics, prebiotics, and fecal microbiota transplantation, as well as their impact on metabolic health and weight management. Finally, the article highlights new research directions, such as personalized microbiome therapies, gut-brain interactions, and multi-omics approaches, suggesting that these could lead to more effective treatments for obesity. The article concludes by emphasizing the importance of further research into the gut microbiome to develop targeted interventions to combat obesity.

Keywords: obesity, gut microbiota, gene.

1. Introduction

Obesity is now considered a critical community health issue worldwide, the prevalence of obesity has increased significantly. The World Health Organization (WHO) reports that the global obesity rate has tripled since 1975. In 2016, more than 1.9 billion adults were considered overweight, with over 650 million falling into the obese category. The rapid increase in obesity rates has not only caused significant health problems but also raised medical costs related to comorbidities such as cardiovascular diseases, diabetes, and certain types of cancer.

Obesity results from the complex interplay of several factors, including genetics, environment, and lifestyle. Recent research indicates that the gut microbiome significantly influences the onset of obesity, presenting novel possibilities for treatment. The significance of the gut microbiome in regulating metabolism, energy storage, and inflammatory responses provides a solid framework for investigating its impact on obesity. This article aims to analyze the significance of the gut microbiome in obesity and explore the potential of regulating the microbiome as a strategy for managing obesity.

2. Gut Microbiota

2.1. Basic Composition of Gut Microbiota

The human gut contains approximately 100 trillion microorganisms, roughly ten times the number of human cells. The two main phyla of bacteria in the gut are Firmicutes and Bacteroidetes, along with

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/75/2024.LA19353

smaller numbers of Actinobacteria, Proteobacteria, and Verrucomicrobia. Although Firmicutes and Bacteroidetes make up the majority of the gut microbiota, their balance can significantly impact metabolism and obesity.

The microbiota in each person's gut can vary greatly, influenced by diet, age, genetics, and environmental factors. However, dysbiosis refers to an imbalance in the microbial community of the gut composition, is often linked to various metabolic disorders, including obesity.

2.2. Functions of Normal Gut Microbiota

In a healthy state, the gut microbiota performs several important functions:

2.2.1. Metabolism. The gut microbiota helps break down complex carbohydrates that the human body cannot digest on its own. These carbohydrates undergo fermentation to generate short-chain fatty acids (SCFAs), mainly acetate, butyrate, and propionate, all essential for maintaining gut health, metabolism, and energy production. SCFAs have the capability to influence lipid metabolism as well, preventing excess fat from accumulating in the body.

2.2.2. Regulation of the Immune System. The gut microbiome engages with the immune system, assisting in the differentiation between harmful pathogens and beneficial microorganisms. This interaction occurs through the regulation of regulatory T cell activity, which helps maintain immune tolerance and prevent excessive inflammatory responses.

2.2.3. Maintaining Intestinal Barrier Integrity. The gut microbiota supports the integrity of the intestinal epithelial barrier, a crucial layer that prevents harmful pathogens, toxins, and antigens from entering the bloodstream. By encouraging the generation of mucus and tight junction proteins, the microbiota helps maintain a physical barrier to ensure normal gut function and reduce the risk of systemic inflammation.

2.2.4. Immune. Modulation and Inflammation Besides impacting local immune functions, the gut microbiota also influences systemic immune functions. Research has indicated that dysbiosis can result in chronic low-grade inflammation, a key feature of metabolic conditions such as obesity and type 2 diabetes. Gut bacteria interact with the immune system through receptors like Toll-like receptors (TLRs) and NOD-like receptors (NLRs), which recognize bacterial components and modulate immune responses. A healthy gut microbiome can suppress inappropriate inflammatory responses, while dysbiosis may result in immune imbalance, contributing to chronic diseases.

3. Changes in Gut Microbiota in Obesity

3.1. Microbial Diversity and Firmicutes/Bacteroidetes Ratio

One of the most significant changes in the gut microbiota of obese individuals is the reduction in bacterial diversity. In particular, a higher proportion of Firmicutes compared to Bacteroidetes is often observed in both animal models and humans with obesity. This change in ratio is linked to an increase in ability to extract energy from food, as Firmicutes are particularly efficient at metabolizing dietary carbohydrates and extracting calories derived from polysaccharides that would otherwise be indigestible.

For example, Turnbaugh et al. [1] transplanted the gut microbiota of obese mice into germ-free mice and found that the mice that received the treatment gained weight. despite having the same caloric intake as the control group. This experiment demonstrates that gut microbiota can significantly influence fat storage by enhancing energy extraction and utilization.

DOI: 10.54254/2753-8818/75/2024.LA19353

3.2. Gut Microbiota and Fat Accumulation

Short-chain fatty acids (SCFAs) produced by intestinal bacteria, including butyrate, acetate, and propionate, have the ability to impact the storage of fats. SCFAs promote fat accumulation by increasing insulin secretion and altering metabolic pathways that regulate fat storage. Acetate can promote the release of insulin from pancreatic β-cells, promoting fat storage in adipose tissue. Propionate, on the other hand, can reduce fat accumulation by inhibiting cholesterol synthesis in the liver. In obese individuals, a reduction in butyrate-producing bacteria is associated with metabolic changes that tend to promote fat accumulation and decrease insulin sensitivity.

3.3. Impact of Dysbiosis on Host Metabolism

Dysbiosis disturbs important metabolic pathways, resulting in insulin resistance, impaired glucose tolerance, and fat buildup. The gut microbiota impacts bile acid metabolism, which in turn controls the absorption of lipids and the balance of glucose levels in the body. Changes in bile acid metabolism in obese individuals can lead to metabolic disturbances, including impaired lipolysis and increased fat storage. Moreover, dysbiosis can result in a reduction of helpful bacteria like Bifidobacterium, which are linked to enhanced gut barrier function and improved metabolic health.

4. Mechanisms by Which Gut Microbiota Influence Obesity

The relationship between gut microbiota and obesity is rooted in several physiological mechanisms that mediate the interactions between microbial communities and host metabolic pathways. These mechanisms include energy extraction efficiency, regulation of intestinal barrier integrity, impact on systemic inflammation, and effects on insulin sensitivity, lipid metabolism, and hunger signals.

4.1. Energy Extraction Efficiency

One of the earliest findings regarding the relationship between gut microbiota and obesity is the role of gut bacteria in energy extraction from food. Gut bacteria, especially Firmicutes, can efficiently break down complex carbohydrates into SCFAs. This increased ability to extract calories from indigestible polysaccharides enhances the energy available to the host, leading to weight gain. Turnbaugh et al. [2] demonstrated that the microbiota associated with obesity individuals has a higher capacity for energy harvesting, leading to fat accumulation even in the absence of excessive caloric intake.

4.2. Intestinal Barrier Function and Inflammation

Obesity is often associated with impaired intestinal barrier function, allowing bacterial endotoxins, particularly lipopolysaccharides (LPS), to enter the bloodstream, leading to metabolic endotoxemia. This process is crucial because LPS induces systemic inflammation, a significant contributor to the development of insulin resistance and metabolic issues dysfunction. Cani et al. [3] found that a high-fat diet disrupts the gut microbiota, leading to increased intestinal permeability and elevated LPS levels in the blood. Once LPS enters the bloodstream, it activates immune responses via Toll-like receptors, triggering chronic low-grade inflammation. This inflammation, in turn, promotes the emergence of insulin resistance, which serves as a precursor to metabolic syndrome and obesity.

4.3. Insulin Sensitivity and Glucose Homeostasis

An important mechanism linking gut microbiota and obesity is its effect on insulin sensitivity and glucose metabolism. Certain bacterial species, such as Bifidobacterium, have been shown to enhance insulin sensitivity by enhancing intestinal barrier integrity and decreasing overall inflammation in the body. Studies have found that individuals with higher levels of Bifidobacterium exhibit better glucose regulation and lower levels of inflammation.

Conversely, dysbiosis reduces the abundance of these beneficial bacteria, impairing glucose metabolism and contributing to the development of insulin resistance. Restoring a healthy gut microbiota through dietary changes or supplementation with specific bacterial strains can help improve insulin sensitivity and glucose control.

DOI: 10.54254/2753-8818/75/2024.LA19353

4.4. SCFAs and Lipid Metabolism

As previously mentioned, SCFAs are essential to lipid metabolism. The gut bacteria produces butyrate, acetate, and propionate, regulate the gene expression involved in fat storage and oxidation. Butyrate is particularly important to preserve the integrity of the intestinal lining and regulating inflammatory responses, both of which are critical for metabolic health.

SCFAs also regulate energy balance by acting on G-protein-coupled receptors (GPCRs), which balance energy by modulating insulin sensitivity and lipid metabolism. Dysbiosis leads to reduced SCFA production, resulting in a metabolic imbalance that favors fat accumulation and obesity.

4.5. Gut-Brain Axis

The gut-brain axis describes the two-way communication between the digestive system and the central nervous system. Through hormonal, neural, and immune pathways, gut microbiota can influence various behavioral aspects, including appetite, cravings, and emotional eating.

The gut microbiota has the ability to generate neurotransmitters such as serotonin and gamma-aminobutyric acid (GABA), which are essential for regulating mood and responding to stress. An imbalance in the microbiota, known as dysbiosis, is linked to alterations in neurotransmitter production and disturbances in the gut-brain communication pathway, leading to unhealthy eating behaviors and weight gain.

4.6. Influence on Hormones and Satiety

Gut microbiota influence the secretion of hormones such as leptin, ghrelin, and peptide YY, which are involved in regulating hunger and satiety signals. Ghrelin is known as the “hunger hormone”, while leptin is referred to as the “satiety hormone”. Dysbiosis can disrupt these hormonal signals, leading to increased feelings of hunger or reduced satiety, driving overeating and contributing to obesity.

Research by Everard et al. [4] showed that restoring the balance of gut microbiota through dietary interventions or probiotic supplementation can help regulate these hormonal pathways, possibly lowering the risk of overeating and promoting weight loss.

5. Factors Affecting Gut Microbiota and Obesity Regulation

Several external factors can influence the composition and function of the gut microbiota and thus affect obesity outcomes. These factors comprise diet, medication, lifestyle, and genetic predisposition.

5.1. Influence of Diet

Diet plays a crucial role in shaping the composition of the gut microbiota. Diets high in saturated fats and simple sugars promote the growth of Firmicutes while reducing the proportion of Bacteroidetes. This modification is linked to elevated fat accumulation and obesity. In contrast, a diet rich in dietary fiber supports the increase of helpful bacteria like bifidobacteria and lactobacilli, which promote metabolic health and prevent obesity [5].

A fiber-rich diet can also enhance the production of SCFAs, which have anti-inflammatory properties and help regulate lipid metabolism. Studies show that individuals who consume diets rich in whole grains, fruits, and vegetables tend to have a more diverse gut microbiota, which is associated with better metabolic outcomes and a lower risk of obesity.

5.2. Antibiotics and Medications

Antibiotics significantly influence the gut microbiota by destabilizing the balance of microbial populations. Using antibiotics is important for curing bacterial infections, but excessive use can cause lasting alterations in the gut microbiota. Thuny et al. [6] showed that prolonged use of antibiotics is associated with weight gain, possibly due to the reduction of beneficial bacteria and the overgrowth of opportunistic pathogens.

Additional medications, like Proton pump inhibitors (PPIs) and nonsteroidal anti-inflammatory medications (NSAIDs), can similarly impact the microbiota by changing the gut environment,

DOI: 10.54254/2753-8818/75/2024.LA19353

increasing the likelihood of dysbiosis. The long-term effects of these medications on weight and metabolic health remain a topic of ongoing research.

5.3. Lifestyle Factors

Lifestyle factors such as physical activity, sleep, and stress play an important role in shaping the composition of the gut microbiota. Regularly exercise has been shown to increase the diversity of the Gut microbiota can be enhanced, especially by boosting the population of beneficial bacteria like bifidobacteria, which are linked to improved metabolic results.

On the other hand, chronic stress can have an adverse effect on the gut microbiome, leading to dysbiosis and increased gut permeability. Stress-related alterations in the gut microbiome are believed to be linked to the development of metabolic disorders, including obesity. Thus, sufficient sleep and stress management are crucial for maintaining a healthy gut microbiota.

5.4. Gene-Environment Interactions

The makeup of the gut microbiota is shaped by both environmental influences and the genetic makeup of the host. Genetic factors can impact how the gut microbiota is structured and how it functions, thereby influencing metabolic health. Two studies have shown that even in genetically identical individuals, differences in the gut microbiota can lead to different obesity outcomes, depending on diet, lifestyle, and environmental exposures [7].

6. The Potential of Gut Microbiota in Obesity Interventions

Targeting the gut microbiota offers a promising therapeutic approach for treating obesity. Various interventions, including probiotics, prebiotics, dietary changes, fecal microbiota transplantation (FMT) has demonstrated promise in restoring microbial balance and enhancing metabolic health.

6.1. Probiotic and Prebiotic Interventions

Probiotics are active microorganisms that provide health benefits to the host when consumed sufficient quantities. Some probiotic strains, specifically those from the Lactobacillus and Bifidobacterium genera, have demonstrated the ability to control fat metabolism, decrease inflammation, and support weight loss. Prebiotics are dietary fibers that nourish beneficial bacteria and have garnered attention for their ability to enhance gut health and support weight management.

A study by Kadooka et al [8]. showed that supplementation with Lactobacillus gasseri significantly reduced abdominal fat, suggesting that specific probiotic strains may help reduce visceral obesity. Prebiotics are particularly effective because they not only provide beneficial bacteria but also supply the necessary nutrients for their growth, creating an environment conducive to gut health.

6.2. Microbiota-Based Drugs and Therapies

Emerging research indicates that microbial byproducts, including short-chain fatty acids (SCFAs) and bile acids, may act as possible targets for obesity therapy. Bifidobacterium, known for maintaining gut barrier integrity, shows promise in reducing inflammation and improving metabolic outcomes in obese individuals.

Preclinical studies have also explored the possibility of using microbiota-based drugs to modulate gut microbial communities and treat obesity. These drugs could target specific microbial pathways involved in energy metabolism, inflammation, and fat storage. One potential approach to managing obesity and its associated health issues is through therapies that target the stimulation of helpful microbial metabolites like butyrate, while also minimizing harmful metabolites such as lipopolysaccharides.

6.3. Fecal Microbiota Transplantation

Fecal microbiota transplantation (FMT) entails the transfer of gut microbiota from a healthy donor to a recipient to restore microbial balance. FMT has been successful in treating Clostridium difficile

DOI: 10.54254/2753-8818/75/2024.LA19353

infections and is now being explored as a potential intervention for metabolic disorders, including obesity.

In obesity research, Ridaura et al. [9] demonstrated that transplanting the gut microbiota from lean individuals into obese mice improved insulin sensitivity and metabolic outcomes. Similar human studies have also shown positive results, suggesting that FMT may be an effective strategy for reshaping the gut microbiota and reversing obesity-related metabolic disorders.

However, FMT presents challenges, including the risk of transferring harmful pathogens while transplanting beneficial bacteria. Moreover, the long-term impacts of FMT on gut microbiota and metabolic well-being are still not clearly understood. While FMT shows promise, further research is needed to ensure its safety and efficacy in treating obesity.

7. Future Research Directions

As we gain a better comprehension of gut microbiota and its influence on obesity, several research areas are emerging that may lead to more effective interventions, including personalized medicine, the gut-brain axis, and multi-omics approaches.

7.1. Personalized Medicine and Gut Microbiota Therapies

Personalized medicine seeks to customize treatment strategies according to individual patient characteristics of each patient, a concept particularly relevant in the field of gut microbiota. Advances in microbiome sequencing technology have made it possible to analyze the composition of an individual's gut microbiota, providing insights into how it affects metabolic health.

By identifying specific microbial imbalances or deficiencies, clinicians can design personalized dietary interventions or probiotic supplementation plans targeting an individual's unique microbiome characteristics. Precision nutrition, which adjusts dietary recommendations based on an individual's gut microbiota, has the potential to optimize weight management and improve metabolic outcomes.

7.2. Gut-Brain Microbiota Interactions

The gut-brain axis is becoming a more significant focus in obesity research, highlighting the intricate relationships between gut microbiota the central nervous system, and behavior. Future studies should focus on how gut bacteria influence cognitive and emotional processes related to eating behaviors, such as stress-induced overeating, food cravings, and emotional regulation.

Additionally, research on psychobiotics--probiotics that influence the gut-brain axis--may offer new therapeutic options for addressing the psychological aspects of obesity. By modulating the gut microbiota to improve mood and reduce stress, psychobiotics could help alleviate behaviors that lead to overeating and weight gain.

7.3. Multi-Omics Approaches

Multi-omics strategies combine data from genomics, transcriptomics, proteomics, and metabolomics, offering valuable insights into the intricate relationships between gut microbiota and host metabolism. These methods enable researchers to explore how gut microbes affect host gene expression, protein production, and metabolic pathways [10].

For example, metabolomics can analyze microbial metabolites, such as SCFAs and bile acids, which play critical roles in regulating host metabolism. By combining multi-omics data with microbiome analysis, researchers can gain a more comprehensive understanding of how gut bacteria promote obesity at the molecular level.

8. Conclusion

The connection between obesity and gut microbiota is an expanding area of research, with growing evidence indicating that gut microbiota plays a crucial role in regulating metabolic health. Dysbiosis is associated with various mechanisms of obesity, including enhanced energy extraction from food, impaired intestinal barrier function, systemic inflammation, and disruptions in appetite and satiety

DOI: 10.54254/2753-8818/75/2024.LA19353

hormone regulation. Research on the gut microbiota opens new avenues for obesity treatment, ranging from probiotics and prebiotic interventions to fecal microbiota transplantation and microbiota-based drugs. These therapies hold promise for restoring microbial balance, improving metabolic outcomes, and providing more personalized solutions for weight management. Dietary interventions, particularly those promoting fiber-rich, microbiota-friendly diets, remain foundational for gut health and obesity prevention. At the same time, the potential of personalized medicine based on individual gut microbiota characteristics could revolutionize our approach to obesity treatment, offering tailored solutions for specific microbial imbalances that contribute to weight gain. Future research will continue to explore the complex relationship between gut microbiota and host metabolism, with particular attention to the gut-brain axis, gene-environment interactions, and multi-omics approaches .As we deepen our comprehension of the gut microbiome, we are getting closer to developing more effective and sustainable solutions to address the global obesity epidemic.

References

[1] Ley, R. E. , Backhed, F. , Turnbaugh, P. , Lozupone, C. A. , Knight, R. D. , & Gordon, J. I. . (2005). Obesity alters gut microbial ecology. *Proceedings of the National Academy of Sciences of the United States of America, 102*(31), 11070-5. https://doi.org/10.1073/pnas.0504978102

[2] Turnbaugh, P. J. , Ley, R. E. , Mahowald, M. A. , Magrini, V. , Mardis, E. R. , & Gordon, J. I. . (2006). An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature, 444*(7122), 1027-1031. https://doi.org/10.1038/nature05414

[3] Cani, P. D. , Possemiers, S. , Wiele, T. V. D. , Guiot, Y. , Everard, A. , & Rottier, O. , et al. (2018). Changes in gut microbiota control inflammation in obese mice through a mechanism involving glp-2-driven improvement of gut permeability. *Gut. 58*(8):1091-103. https://doi.org/10.2337/db06-1493.

[4] Everard, A. , & Cani, P. D. (2013). The gut microbiota as a significant contributor to metabolic disorders. *Nature Reviews Gastroenterology & Hepatology, 11*, 55–67. https://doi.org/10.1038/nrgastro.2013.151

[5] Turnbaugh, P. J. , B Ckhed, F. , Fulton, L. , & Gordon, J. I. . (2008). Dietd obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host & Microbe, 3*(4), 213-223. https://doi.org/10.1038/nature06244

[6] Thuny, F. , Di, S. G. , Belliard, O. , Avierinos, J. F. , Pergola, V. , & Rosenberg, V. , et al. (2005). Risk of embolism and death in infective endocarditis: prognostic value of echocardiography: a prospective multicenter study. *Circulation, 14*(11), 31. https://doi.org/10.1161/CIRCULATIONAHA.105.584522

[7] Backhed, F. , Ding, H. , Wang, T. , Hooper, L. V. , Koh, G. Y. , & Nagy, A. , et al. (2004). The gut microbiota as an environmental that regulates fat storage. *Proceedings of the National Academy of Sciences, 101*(44), 15718-15723. https://doi.org/10.1073/pnas.0407076101.

[8] Kadooka, Y., Sato, M., Imaizumi, K., Ogawa, A., Ikuyama, K., Akai, Y., Okano, M., & Tsuchida, T. (2010). Regulation of abdominal adiposity by probiotics (Lactobacillus gasseri SBT2055) in adults 224455555with obese tendencies in a randomized controlled trial. *European Journal of Clinical Nutrition, 4*(6), 636-643. https://doi.org/10.1038/ejcn.2010.19.

[9] Turnbaugh, P. J. , Ridaura, V. K. , Faith, J. J. , Rey, F. E. , Knight, R. , & Gordon, J. I. . (2009). The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice. *science translational medicine, 1*(6), 6-6. https://doi.org/10.1126/science.1241214.

[10] Suravajhala, P. , Kogelman, L. J. A. , & Kadarmideen, H. N. . (2016). Multi-omic data integration and analysis using systems genomics approaches: methods and applications in animal production, health and welfare. *Genetics Selection Evolution*, 48(1), 38. https://doi.org/10.1186/s12711-016-0217-x.