Proceedings of ICBioMed 2025 Symposium: AI for Healthcare: Advanced Medical Data Analytics and Smart Rehabilitation
DOI: 10.54254/2753-8818/2025.AU23391

A New Perspective on Exercise Rehabilitation Strategies Based on Hemodynamic Optimization: Enhancing Skeletal Muscle Repair

Yuanshu Lu

Shanghai World Foreign Language Academy, Shanghai, China
luyuanshu_jenny@163.com

Abstract: The integration of hemodynamics into exercise rehabilitation has gained attention as a promising approach to enhance skeletal muscle repair. This study connects the physiological linkage between hemodynamics and muscle regeneration, focusing on how exercise-induced hemodynamic changes can optimize recovery outcomes. By reviewing recent literature, the research examines the mechanisms of blood flow restriction (BFR) training and its impact on muscle repair, alongside other emerging techniques like compression therapy. The key research question is how hemodynamic parameters can be adapted to design personalized rehabilitation strategies. Findings suggest that optimizing blood flow shear stress and oxygenation levels could improve muscle regeneration efficiency, particularly for sports injuries. This study proposes a theoretical framework for tailoring exercise prescriptions based on hemodynamic data, offering a low-cost, non-invasive solution. However, challenges such as accessibility and patient compliance remain. This work emphasizes the potential of hemodynamic-focused rehabilitation and calls for further exploration into integrating wearable technology for real-time monitoring, contributing to precision rehabilitation in sports science and therapy.

Keywords: Hemodynamics, Skeletal Muscle Repair, Exercise Rehabilitation, Blood Flow Restriction, Personalized Therapy

1. Introduction

Skeletal muscle injuries are common in sports and daily activities, often requiring effective rehabilitation to restore function. Recent researches in sports science have highlighted the critical role of hemodynamics—the study of blood flow dynamics—in muscle repair processes [1]. Exercise, as a non-invasive intervention, influences blood flow distribution, oxygenation, and nutrient delivery, all of which are essential for regeneration [2]. However, traditional rehabilitation approaches often overlook individual hemodynamic variations, limiting their efficacy. This paper investigates how exercise-induced hemodynamic changes can be utilized to enhance skeletal muscle repair, with a focus on emerging techniques like blood flow restriction (BFR) training. BFR, which modulates blood flow using controlled pressure, has shown promise in accelerating recovery by enhancing muscle protein synthesis and vascular adaptation [3]. The research question driving this study is: how can hemodynamic parameters be optimized to design personalized rehabilitation strategies? By synthesizing current literature, this study employs a theoretical approach, analyzing mechanisms such

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.AU23391

as shear stress and angiogenesis in muscle repair [4]. The significance of this work lies in its potential to shift rehabilitation paradigms toward precision medicine, offering tailored solutions for athletes and patients. While experimental validation is beyond the scope due to time constraints, the proposed framework leverages existing data to suggest practical applications. This article aims to bridge the gap between hemodynamic research and clinical practice, providing a foundation for future innovations in exercise-based rehabilitation within the fields of sports science and physical therapy.

2. The role of hemodynamics in exercise rehabilitation

2.1. Physiological link between blood flow and muscle repair

Blood flow is a key driver in skeletal muscle repair, delivering oxygen, nutrients, and growth factors while removing metabolic byproducts, especially after exercise-induced injury [2]. Increased circulation post-exercise activates satellite cells, promoting their proliferation and differentiation into myotubes to repair damaged fibers by enhancing nutrient and signaling molecule availability [2]. Blood flow also influences the microvascular environment via shear stress, which upregulates vascular endothelial growth factor (VEGF), driving capillary proliferation. Milkiewicz et al. showed that elevated shear stress in a rat model (via chronic stimulation or prazosin) increased the capillary-to-fiber ratio by 40% over 14 days [4]. Additionally, Hudlicka and Brown noted that shear stress triggers nitric oxide (NO) release, enhancing endothelial function and vascular remodeling to support oxygen delivery and metabolic adaptation [1, 5]. In ischemic conditions, VEGF rises but fails to drive capillary growth without restored flow, highlighting blood flow's active role in regulating repair through mechanical and molecular pathways [1].

2.2. Mechanisms of exercise-induced hemodynamic regulation

Exercise impacts hemodynamic parameters like heart rate, blood pressure, and shear stress, influencing skeletal muscle repair [6]. Tidball noted that eccentric contractions increase local blood flow and shear stress, disrupting myofibrillar structures and triggering repair via blood-borne inflammatory cells and growth factors, with shear stress boosting VEGF expression and capillary growth [7]. Karlsdottir et al. found that moderate-intensity resistance exercise (60%-70% of 1RM) raises diastolic and mean arterial pressures without affecting left ventricular function, supporting muscle perfusion across healthy individuals, coronary artery disease patients, and those with congestive heart failure [6]. Hudlicka and Brown showed that electrical stimulation, mimicking exercise, restores arteriolar dilatation in ischemic muscles, increasing shear stress and VEGF receptor 2 expression, leading to enhanced capillary growth compared to untreated ischemic states [1]. Aerobic training improves systemic oxygenation, while resistance exercise targets localized perfusion, highlighting exercise as a key regulator of blood flow dynamics for optimizing muscle repair in rehabilitation [6].

3. Current blood flow optimization techniques in rehabilitation

3.1. Principles and applications of blood flow restriction (BFR) training

Blood flow restriction (BFR) training enhances skeletal muscle repair in rehabilitation by applying external pressure (e.g., cuffs) to partially restrict venous return while maintaining arterial inflow, creating a hypoxic environment that increases metabolic stress [2, 3]. Damas et al. showed that this stress boosts muscle protein synthesis (MPS) via anabolic signaling at low intensities (20%-30% 1RM), ideal for patients recovering from injuries or surgery [3]. Clifford et al. found BFR reduced soreness and improved functional recovery (e.g., jump height) in a drop-jump injury model, due to

DOI: 10.54254/2753-8818/2025.AU23391

enhanced microvascular perfusion and ECM remodeling [5]. Karlsdottir et al. confirmed BFR's safety, noting moderate-intensity exercise (60%-70% 1RM) raises arterial pressures without affecting left ventricular function, even in cardiovascular patients [6]. Mechanistically, BFR upregulates VEGF and IGF-1, promoting shear stress-induced angiogenesis, making it a practical, low-load tool for early-stage rehabilitation [1].

3.2. Advances in complementary methods

Compression therapy, a passive complement to BFR, optimizes hemodynamics in rehabilitation by applying graduated pressure to limbs, enhancing venous return, reducing edema, and improving microvascular circulation [5]. Clifford et al. showed that post-exercise compression garments, paired with collagen peptide supplementation, accelerated ECM remodeling and functional recovery in a drop-jump injury cohort by improving lymphatic drainage and reducing inflammation, supporting satellite cell activity [2, 5]. Hudlicka and Brown found that chronic electrical stimulation with compression-like effects in ischemic muscles restored endothelial function and increased capillary surface area, enhancing oxygen delivery in peripheral vascular disease patients [1]. Innovations like dynamic compression devices, which adjust pressure in real-time, further refine its application by targeting venous flow velocity [7]. Compared to BFR, compression therapy is less resource-intensive and ideal for later recovery, providing a versatile, blood flow-focused toolkit for rehabilitation across diverse patient needs [5].

4. Designing rehabilitation strategies based on hemodynamics

4.1. Selection of key hemodynamic parameters

Personalized rehabilitation strategies rely on hemodynamic parameters like shear stress and oxygenation to enhance skeletal muscle repair [4]. Shear stress, the frictional force of blood flow on the endothelium, upregulates VEGF and VEGFR-2, promoting angiogenesis. Milkiewicz et al. showed that prazosin in rats quadrupled capillary shear stress in 3-4 days, tripling PCNA-positive capillaries and VEGF expression [4]. Hudlicka and Brown found chronic stimulation of ischemic muscles increased shear stress, restoring capillary growth [1]. Oxygenation supports metabolic efficiency during repair; Gnimassou linked it to improved mitochondrial function and satellite cell proliferation [5]. Nachman et al. validated non-invasive oxygenation monitoring via photoplethysmography (PPG) in a swine model ($r=0.89$-0.93 vs. invasive measures), enabling real-time assessment [7]. Other parameters like heart rate, stable across healthy and cardiac patients during resistance exercise, provide additional context [6]. Integrating shear stress and oxygenation allows tailored rehabilitation using wearable devices for precision [7].

4.2. Theoretical framework for personalized exercise prescriptions

A theoretical framework for personalized exercise prescriptions based on hemodynamics integrates individual blood flow profiles to optimize muscle repair outcomes [2]. This approach begins with assessing baseline hemodynamic parameters, such as shear stress, oxygenation, and blood pressure—using non-invasive technologies like PPG devices, which Nachman et al. validated for continuous monitoring with high intraclass correlations (e.g., 0.92 for systolic blood pressure) [7]. For patients exhibiting high shear stress responses, blood flow restriction (BFR) training is ideal; Dumont et al. noted that BFR enhances satellite cell activation and protein synthesis under hypoxic conditions, accelerating repair without excessive mechanical load [2]. Conversely, individuals with low oxygenation levels, as Gnimassou suggests, may benefit from aerobic exercise to boost systemic oxygen delivery and metabolic recovery [5].

DOI: 10.54254/2753-8818/2025.AU23391

This framework also incorporates complementary interventions. Clifford et al. found that collagen peptide supplementation paired with BFR or compression therapy amplified extracellular matrix (ECM) remodeling, reducing recovery time post-injury [5]. For example, a patient with elevated shear stress and moderate oxygenation might receive a hybrid prescription: low-intensity BFR (20%-30% 1RM) thrice weekly, supplemented with collagen peptides, to target both vascular and structural repair [1,5]. Adjustments are guided by real-time data—e.g., if cardiac output drops (monitored via PPG) [7], intensity could shift to aerobic intervals. This adaptability contrasts with one-size-fits-all protocols, addressing individual variability in hemodynamic responses noted by Karlsdottir et al. [6]. While theoretical, this framework leverages existing evidence to propose a dynamic, patient-centered model, setting the stage for empirical validation and broader clinical application in sports rehabilitation.

5. Potential and challenges of strategy implementation

5.1. Enhancing muscle repair efficiency

In exercise rehabilitation, enhancing muscle repair efficiency is a crucial objective. Optimizing hemodynamics plays a key role in reducing recovery time and improving functional recovery [2]. It has been shown that improved blood flow facilitates the delivery of oxygen and nutrients to damaged muscle tissues, accelerating the repair and regeneration process [2]. By optimizing circulation, metabolic waste products are more rapidly removed, which not only reduces cell damage but also prevents the escalation of inflammation [5]. This process contributes to the overall recovery of muscle tissues and enhances the patient's ability to perform physical tasks.

Electrical stimulation has shown significant potential in improving blood flow to ischemic muscles [1]. By mimicking neural signals, electrical stimulation activates damaged muscles, leading to vasodilation and an increase in blood flow [1]. This enhances the delivery of oxygen and nutrients to the tissue, while also promoting healing of the injured area. This is particularly important in muscles that have been subjected to prolonged ischemia or poor circulation, as electrical stimulation can improve microcirculation and promote blood flow, thereby reducing muscle pain and discomfort and accelerating recovery [1].

In clinical settings, this strategy holds considerable promise, particularly in post-injury or post-surgical recovery. Electrical stimulation and blood flow optimization provide an effective rehabilitation approach for patients [8]. By carefully controlling the frequency and intensity of electrical stimulation, it is possible to achieve optimal blood flow improvement without causing adverse effects, thus enhancing the rate and quality of muscle repair [1]. Despite the significant potential of this strategy, there are still challenges to overcome. One key issue is how to fine-tune the parameters of electrical stimulation to maximize blood flow and muscle repair efficiency.

Additionally, due to the individual differences among patients, personalized treatment plans remain a challenge. Different patients have varying levels of muscle damage and recovery needs, making it necessary to tailor rehabilitation programs to specific circumstances, which requires further research and clinical practice.

5.2. Issues of accessibility and patient compliance

The accessibility of blood flow restriction (BFR) equipment is often hindered by the high costs associated with acquiring the necessary devices and the training required for both healthcare professionals and patients [3]. While BFR therapy has been shown to enhance muscle repair and strength [3], the substantial financial investment in specialized equipment presents a barrier to its widespread implementation, particularly in resource-limited settings. Moreover, the need for proper training and supervision to ensure the safe and effective use of BFR devices further complicates their

DOI: 10.54254/2753-8818/2025.AU23391

availability and increases the overall cost of treatment. As a result, despite its potential benefits, the adoption of BFR therapy in clinical practice may be restricted by these financial and educational constraints.

Patient adherence to long-term rehabilitation protocols remains a significant challenge, especially when such protocols are complex or require frequent visits to specialized healthcare facilities. Consistent compliance with exercise regimens and treatment schedules is critical for achieving optimal recovery outcomes. However, the burden of regular appointments and the need for continued supervision often lead to decreased adherence over time. To address this issue, cost-effective solutions such as wearable devices have emerged as a promising alternative [7]. Wearables, which allow for continuous monitoring and management of exercise protocols, offer greater convenience and flexibility for patients, enabling them to adhere to their rehabilitation plans more easily [7]. These devices can provide real-time feedback and track progress, thus improving motivation and encouraging consistent participation. Moreover, their portability and ease of use make them an attractive option for patients who may face challenges with traditional, clinic-based rehabilitation approaches. However, the adoption of wearable devices in BFR therapy also raises concerns regarding their effectiveness, accuracy, and long-term feasibility, requiring further investigation to determine how best to integrate them into comprehensive rehabilitation strategies [7].

6. Discussion

Hemodynamics plays a complementary role in traditional rehabilitation strategies by focusing on the dynamic aspects of blood flow [1]. By optimizing circulation, it enhances the delivery of essential nutrients and oxygen to damaged tissues, accelerates metabolic waste removal, and fosters a more conducive environment for muscle repair [2]. This approach aligns well with traditional rehabilitation methods, which typically emphasize mechanical and physiological recovery through exercise and physical therapy. However, hemodynamic optimization goes a step further by addressing the vascular side of recovery, which can be particularly beneficial in cases of ischemic or impaired circulation [1].

While promising, the immediate application of hemodynamic optimization in clinical rehabilitation is limited by a lack of empirical data. While theoretical frameworks and early studies suggest its potential benefits, the current body of research is insufficient to provide definitive evidence on its efficacy across different patient populations and injury types [4]. Much of the existing data comes from experimental settings or small-scale trials, making it difficult to generalize findings to larger clinical practices. The absence of long-term, large-scale studies further limits our understanding of the true impact of hemodynamic optimization on muscle recovery and functional outcomes [1].

Therefore, while hemodynamics presents an exciting avenue for enhancing rehabilitation strategies, its widespread application hinges on future validation through rigorous research. Larger clinical trials, longitudinal studies, and more robust datasets are necessary to establish the most effective protocols for integrating blood flow optimization into rehabilitation plans [8]. Additionally, understanding the specific mechanisms by which hemodynamic interventions influence muscle repair will be crucial in developing standardized, evidence-based treatment protocols that can be widely adopted in clinical settings [9]. Without such empirical backing, the integration of hemodynamic optimization into mainstream rehabilitation practices remains speculative.

7. Conclusion

Wearable photoplethysmogram (PPG) devices offer significant potential for real-time monitoring of hemodynamic parameters, such as blood flow and oxygen saturation, during rehabilitation [7]. These devices enable continuous tracking of vascular health, providing clinicians and patients with valuable

DOI: 10.54254/2753-8818/2025.AU23391

insights into the body's response to exercise and rehabilitation protocols [7]. The integration of wearable PPG technology in hemodynamic monitoring can help facilitate dynamic adjustments to rehabilitation plans, ensuring that interventions are tailored to the patient's current physiological state [7]. By tracking blood flow, heart rate, and other hemodynamic markers, these devices allow for immediate feedback, enabling modifications to the intensity, frequency, or duration of exercises based on real-time data [6].

The use of wearable PPG devices could enhance patient adherence to rehabilitation protocols by making monitoring more accessible and convenient [7]. Instead of requiring frequent visits to clinics for assessments, patients can receive continuous monitoring at home, reducing the burden of travel and providing a more personalized approach to rehabilitation. This also allows for better tracking of progress over time, which can be particularly useful for patients who require long-term rehabilitation or those with chronic conditions [7].

Future studies should explore the potential of wearable PPG devices across diverse populations, including individuals with varying levels of physical fitness, age, and medical conditions. The adaptability of these devices in different contexts, such as post-surgery, athletic recovery, or chronic disease management—should be further investigated [7]. Additionally, it will be important to evaluate the accuracy and reliability of wearable devices in capturing hemodynamic parameters across different environments and conditions [7]. Exploring how these devices interact with other rehabilitation technologies, such as electrical stimulation or blood flow restriction, could provide valuable insights into optimizing combined therapies [1,3].

References

[1] *Hudlicka, Olga, and Margaret D. Brown. Adaptation of Skeletal Muscle Microvasculature to Increased or Decreased Blood Flow: Role of Shear Stress, Nitric Oxide and Vascular Endothelial Growth Factor. Journal of Vascular Research, vol. 46, no. 5, 2009, pp. 504-512.*

[2] *Dumont, Nicolas A., et al. Satellite Cells and Skeletal Muscle Regeneration. Comprehensive Physiology, doi:10.1002/cphy.c140068.*

[3] *Damas, Felipe, et al. A Review of Resistance Training-Induced Changes in Skeletal Muscle Protein Synthesis and Their Contribution to Hypertrophy. Sports Medicine.*

[4] *Milkiewicz, Malgorzata, et al. Association Between Shear Stress, Angiogenesis, and VEGF in Skeletal Muscles in Vivo. Microcirculation, vol. 8, no. 4, 2001, pp. 229-241.*

[5] *Hernandez-Torres, F., et al. Macrophages in the Context of Muscle Regeneration and Duchenne Muscular Dystrophy. International Journal of Molecular Sciences, vol. 25, no. 19, 2024.*

[6] *Karlsdottir, Arna E., et al. Hemodynamic Responses Dring Aerobic and Resistance Exercise. Journal of Cardiopulmonary Rehabilitation and Prevention, vol. 22, no. 3, 2002, pp. 170-177.*

[7] *Nachman, Dean, et al. Wireless, Non-invasive, Wearable Device for Continuous Remote Monitoring of Hemodynamic Parameters in a Swine Model of Controlled Hemorrhagic Shock. Scientific Reports, vol. 10, no. 1, 2020, p. 17684.*

[8] *Esteki, Fatemeh Ghashghaei, et al. Exercise-based Cardiac Rehabilitation Improves Hemodynamic Responses After Coronary Artery Bypass Graft Surgery. ARYA Atherosclerosis, vol. 7, no. 4, 2012, pp. 151.*

[9] *Tidball, James G. Mechanisms of muscle injury, repair, and regeneration. Comprehensive Phy siology vol. 1,4 (2011): 2029-62. doi:10.1002/cphy.c1000*