Proceedings of CONF-MPCS 2025 Symposium: Leveraging EVs and Machine Learning for Sustainable Energy Demand Management
DOI: 10.54254/2753-8818/2025.GL24074

Microscopic Interpretation and Macroscopic Applications of the Second Law of Thermodynamics

Yiming Lu

Hebei Normal University, Shijiazhuang, China
LUYIMING9963@163.com

Abstract: The second law of thermodynamics, a fundamental principle of nature, reveals the unidirectional evolution of isolated systems toward equilibrium through the concept of entropy increase. This study explores the microscopic essence of the law from a statistical mechanics perspective, demonstrating that systems naturally evolve toward states of maximum probability. It also examines how this principle manifests across different scales. By analyzing the derivation of entropy production criteria, the study unveils the physical connotation of entropy and its role in equilibrium and non-equilibrium systems. On the macroscopic level, it investigates how entropy increase limits energy conversion efficiency and determines the direction of thermodynamic processes. Case studies on heat engine efficiency and energy conversion mechanisms demonstrate the law's guiding significance for optimizing engineering thermodynamic systems. This research not only deepens the comprehension of the unity between microscopic mechanisms and macroscopic manifestations of the second law, but also provides theoretical support for enhancing energy utilization efficiency and thermodynamic cycle system performance.

Keywords: The second law of thermodynamics, entropy criterion, Clausius inequality, Boltzmann, heat engine

1. Introduction

As a cornerstone of classical thermodynamics, the second law of thermodynamics reveals the inherent irreversibility of energy conversion processes and determines the natural evolution of isolated systems. Its fundamental formulation—the entropy increase principle—states that the total entropy of an isolated system never decreases. This principle describes, on a macroscopic level, the spontaneous trend of systems toward equilibrium, and, on a microscopic level, it reflects the statistical behavior of particle distributions. Since Clausius introduced the macroscopic definition of entropy, Boltzmann further bridged it with macroscopic particle configurations via statistical entropy, providing a quantitative foundation for a probabilistic interpretation of irreversibility. Despite these developments, theoretical challenges remain, particularly regarding the entropy principle's application in non-equilibrium systems and its implications for energy losses and thermodynamic efficiency. This study investigates the second law through its dual microscopic-macroscopic perspectives. Commencing with entropy derivations and the Clausius inequality, the author analyzes how entropy constrains the direction of classical irreversible processes such as heat conduction and free expansion. Furthermore, by applying Carnot's theorem and considering real-world dissipation, the paper analyzes limits of heat engine efficiency and the role of entropy in energy conversion

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.GL24074

system optimization. Through this dual-perspective approach, this research aims to unify statistical interpretations with thermodynamic behavior and provide theoretical guidance for improving energy efficiency in engineering systems.

2. The theoretical framework of the second law of thermodynamics

2.1. Derivation of Clausius' entropy and explanation of the entropy increase principle

The concept of entropy was gradually developed through extensive research, enabling Clausius to resolve the foundation of the second law of thermodynamics for non-cyclic processes. Historically, its this development occurred in two distinct phases. In 1854, Clausius first proposed a mathematical expression of the second law for cyclic processes. About a decade later, he extended the theory to non-cyclic transformations by introducing state functions, based on his microscopic view of matter [1]. The derivation of Clausius entropy begins with his theorem: for any reversible cyclic process, the closed integral of the heat-to-temperature ratio equals zero:

$$\oint {\delta Q_{rev}}/{T} = 0 \tag{1}$$

For irreversible cycles, the integral is less than zero:

$$\oint {\delta Q_{irrev}}/{T} < 0 \tag{2}$$

This leads to the definition of entropy as a state function. Consider a system undergoing a reversible processes from state A to state B via reversible path 1, and returning to A through another reversible path 2, forming a closed cycle. Then:

$$\int_{A\to B}^{1} {\delta Q_{rev}}/{T} + \int_{B\to A}^{2} {\delta Q_{rev}}/{T} = 0 \tag{3}$$

Which implies:

$$\int_{A\to B}^{1} {\delta Q_{rev}}/{T} = \int_{B\to A}^{2} {\delta Q_{rev}}/{T} \tag{4}$$

Thus, the integral value is path-independent, depending solely on the initial and final states. Consequently, there exists a state function \(S \) whose change is given by:

$$\triangle S = S(B) - S(A) = \int_{A}^{B} {\delta Q_{rev}}/{T} \tag{5}$$

This function is entropy. Next, for irreversible systems from A to B followed by a reversible return path, the cyclic integral satisfies:

$$\int_{A\to B}^{irrev} {\delta Q}/{T} + \int_{B\to A}^{rev} {\delta Q_{rev}}/{T} < 0 \tag{6}$$

Given that the integral along the reversible path is -ΔS, substitution leads to

$$\int_{A\to B}^{irrev} {\delta Q}/{T} < S(B)-S(A) \tag{7}$$

DOI: 10.54254/2753-8818/2025.GL24074

For irreversible processes, the integral of the heat-to-temperature ratio is less than the entropy change. In adiabatic processes ($\delta Q = 0$) or isolated systems, this inequality reduces to $\Delta S \geq 0$. The equality holds exclusively for reversible processes, while irreversible processes result in entropy increase.

2.2. Foundation of statistical mechanics: microscopic state probability and the statistical interpretation of entropy

Statistical mechanics explains the physical nature of macroscopic thermodynamic quantities (e.g., entropy) through the motion laws of microscopic particles. In the final decades of the 19th century, Boltzmann interpreted irreversible macroscopic laws, particularly the second law of thermodynamics. The core premise is that macroscopic properties arise from the statistical averaging over a vast number of microscopic states. To clarify this, a distinction must be made with macroscopic and microscopic states [2]. A microscopic state refers to a specific instantaneous configuration of all particles in a system (positions, momenta, etc.), constrained by total energy, volume, and other conditions. For example, the exact positions and velocities of N molecules in a container define a microscopic state. A macroscopic state, on the other hand, describes the system through observable variables like temperature T, pressure P, and volume V. Identical macroscopic states can correspond to the same microscopic state, and the total number of these compatible configurations is referred to as the multiplicity or the number of microstates. The fundamental postulate of statistical mechanics—the assumption of equal a priori probabilities states that for an isolated system (constant energy), all microscopic states satisfying the constraints are equally probable. On this basis, Boltzmann proposed the statistical interpretation of entropy:

$$S=K_B \ln W \tag{8}$$

Physical Interpretation: Entropy quantifies the degree of microscopic states in a system. A larger number of microscopic states corresponds to greater disorder, resulting in a higher entropy value. High-entropy states represent macrostates of higher probability, and systems spontaneously evolve in the direction of entropy increase.

2.3. Thermodynamic characteristics of equilibrium and non-equilibrium systems

An equilibrium state is characterized by uniform macroscopic properties (e.g., temperature, pressure) with no net energy or material flows. According to Callen, "in equilibrium, the entropy of a system attains its maximum value, and all driving forces for irreversible processes (e.g., temperature gradients) vanish" [3]. Equilibrium states satisfy equations of state (e.g., the ideal gas law) and involve reversible processes, such as equal phase transition rates in ice-water coexistence. For isolated systems, equilibrium corresponds to the most probable macrostate in statistical mechanics, characterized by a maximized microstate count Ω.. In contrast, non-equilibrium state exhibits spatiotemporal gradients (e.g., temperature or concentration differences) and irreversible transport processes (e.g., heat conduction, diffusion). As Prigogine emphasized, "the entropy production rate in non-equilibrium systems remains positive, reflecting the directional nature of energy dissipation" [4]. For instance, during heat flow from a hot to a cold region, the system's entropy increases but has not yet reached its maximum value. Non-equilibrium states are further classified into near-equilibrium (linear responses, e.g., Fourier heat conduction) and far-from-equilibrium regimes (nonlinear dissipative structures, e.g., Bénard vortices).

Comparing these states reveals that equilibrium corresponds to maximum entropy in isolated systems, while non-equilibrium systems exhibit continuous entropy increase until equilibrium is attained. To describe such systems, the local equilibrium hypothesis is often employed, assuming that

DOI: 10.54254/2753-8818/2025.GL24074

subsystems within a non-equilibrium system can be approximated as being in internal equlibrium at microscopic scales. In open systems, non-equilibrium steady states (e.g., in living organisms) can be sustained through external energy/matter inputs, though the total entropy (system + environment) still increases.

3. The microscopic origin and equilibrium evolution mechanism of entropy

The Boltzmann entropy formula ($S=k\ln\Omega$) bridges macroscopic entropy with the microscopic state count Ω, unveiling the statistical essence of the second law of thermodynamics: isolated systems spontaneously evolve toward equilibrium states with maximized Ω, corresponding to the most probable distribution of microscopic configurations [5]. For example, during free expansion of a gas, molecular phase-space distributions transition from localized (low Ω) to uniform (high Ω), accompanied by entropy increase. This theory transcends the phenomenological framework of classical thermodynamics by interpreting irreversibility as probabilistic evolution, though its foundational assumption of molecular chaos (Stosszahlansatz) initially sparked debate [6]. The probabilistic nature of microscopic configurations manifests further in equilibrium-state fluctuations. Einstein demonstrated that the second derivative of entropy with respect to macroscopic parameters determines fluctuation magnitudes (e.g., density fluctuations), revealing that microscopic states remain dynamically active even at equilibrium [6]. Boltzmann's framework also underpins quantum statistics: Fermi-Dirac and Bose-Einstein distributions can both be derived as generalized entropy-maximization outcomes under distinct microscopic configuration constraints [5].

The spontaneous evolution of isolated systems toward equilibrium fundamentally arises from the statistical inevitability governing microscopic states. When a system contains a vast number of particles, the microscopic configurations corresponding to equilibrium dominate in phase space with overwhelming probabilistic dominance—the likelihood of uniform distributions far exceeds that of any ordered or localized states. This asymmetry in probability distribution underlies macroscopic irreversibility: although microscopic particles obey time-reversible dynamics, the transition from low-probability non-equilibrium states (e.g., temperature inhomogeneity) to high-probability equilibrium occurs with near certainty, while the reverse process is exponentially suppressed and practically unobservable. Although minor fluctuations persist near equilibrium, their magnitude diminishes rapidly with increasing system size, ensuring the macroscopic state remains statistically stabilized at the entropy-maximized equilibrium. This principle elucidates the microscopic origin of the second law of thermodynamics, revealing that entropy increase fundamentally represents the system's evolution toward higher-probability states.

4. Thermodynamic analysis of macroscopic irreversible processes

The study of entropy changes in irreversible processes reveals the directional nature of system evolution. According to the second law of thermodynamics, entropy in isolated systems undergoing irreversible processes always satisfies $\Delta S > 0$. Clausius asserted: "For naturally occurring irreversible processes, the total entropy of the system and its environment must increase, which fundamentally embodies the thermodynamic arrow of time." In heat conduction, when a high-temperature object (T_1) contacts a low-temperature object (T_2), the emtropy change $Q(1/T_2 - 1/T_1)$ is inherently positive [7]. The entropy gain in the high-temperature object Q/T_1 is always smaller in magnitude than the entropy increase in the low-temperature object Q/T_2. This unidirectional energy transfer, driven by temperature gradients, reflects the irreversible tendency of energy distributions to spontaneously homogenize. Experimental data confirm that entropy production increases with the temperature gradient.

DOI: 10.54254/2753-8818/2025.GL24074

Modern non-equilibrium thermodynamics, developed by the Prigogine school, claims "the essential characteristic of irreversible processes lies in their generation of positive-definite entropy productionσ= ∫ (dS/dt)dV⩾0, which arises from the nonlinear coupling between thermodynamic forces and fluxes within the system" [8]. For example, during the free expansion of a gas, the increased disorder in molecular distribution leads to an entropy increase of approximately Nkln2 (where N is the number of molecules). This irreversible growth in the number of microscopic states confirms the statistical mechanical interpretation of macroscopic entropy changes. If gas in a sealed container is abruptly released into a vacuum, the spatial distribution probability of molecules expands exponentially, with the corresponding jump in configurational entropy directly observable as macroscopic volume expansion. Numerical simulations show that when the number of gas molecules reaches the order of Avogadro's constant, the probability of the system returning to its initial state approaches zero, quantitatively characterizing the irreversible nature of such processes from a probabilistic perspective.

5. Engineering application of the second law of thermodynamics

5.1. Theoretical derivation of the Carnot cycle and heat engine efficiency limits

The Carnot cycle is a theoretical model that defines the operation of an ideal heat engine, with its efficiency limit established by Carnot's theorem. According to Carnot's work, the maximum efficiency of a heat engine depends solely on the temperatures of the high-temperature reservoir (T_H) and the low-temperature reservoir (T_C), expressed as:

$$\eta_{max}=1-\frac{T_C}{T_H} \tag{9}$$

Carnot demonstrated that any heat engine exceeding this efficiency would violate the second law of thermodynamics. He concluded that “the maximum efficiency of a heat engine operating between two given temperatures is independent of the working substance and determined solely by the temperature ratio." This conclusion laid the foundation for the entropic formulation of the second law [9, 10]. Carnot's Theorem reveals a key constraint in energy conversion: only the thermal energy difference between the two reservoirs can be converted into useful work. For example, if T_1 = 500 , andT_2 = 300, the theoretical maximum efficiency is 40%, meaning 60% of the energy must dissipate as waste heat. This efficiency limit resides in the irreversibility of heat transfer, where spontaneous heat flow from hot to cold increase entropy, and this increase cannot be fully eliminated. Practical systems like steam turbines and internal combustion engines typically achieve only 30% –50% of Carnot efficiency due to irreversible factors such as frictional losses and non-equilibrium heat transfer. The Carnot Cycle has four reversible processes: isothermal heat absorption, adiabatic expansion, isothermal heat rejection, and adiabatic compression. During the isothermal expansion phase, the system absorbs heat Q_H from the high-temperature reservoir, causing an increase in entropy $\Delta S_H=\frac{Q_H}{T_H}$. In the isothermal compression phase, the system releases heat Q_C to the low-temperature reservoir, resulting in a decrease in entropy $\Delta S_C=-\frac{Q_C}{T_C}$. Since the total entropy change over the cycle must equal zero, we derive: $\frac{Q_H}{T_H}=\frac{Q_C}{T_C}$. This relationship emphasizes the importance of temperature gradients in determining energy quality. The higher the temperature of the high-temperature reservoir, the greater the proportion of "usable energy" (exergy) contained within its thermal energy.

DOI: 10.54254/2753-8818/2025.GL24074

In practical heat engines, entropy production reduces the efficiency and work output. For example, in gas turbines, a 10% increase in the temperature gradient between the combustion chamber and turbine raises the entropy production rate by about 15%, resulting in an efficiency loss of 2%–3%. The universality of Carnot's theorem extends beyond conventional heat engines. Its efficiency limits apply to emerging energy conversion systems, such as thermoelectric generators and fuel cells. Even at quantum scales, the efficiency of nanoscale heat engines remains bounded by the Carnot efficiency limit, confirming the spatiotemporal universality of Carnot cycle principles.

Carnot's work not only provided the mathematical formulation for the second law of thermodynamics but also inspired Clausius to develop the entropy function. By proving that for reversible cycles

$$\oint {\delta Q}/{T} = 0 \tag{10}$$

while irreversible processes satisfy:

$$\oint {\delta Q}/{T} < 0 \tag{11}$$

the entropic statement of the second law was rigorously established. The principles of the Carnot cycle guide the optimization of real-world energy systems. For instance, in coal-fired power plants, elevating steam temperatures from 300 ℃ to 620℃ can raise Carnot efficiency from 48% to 67%, though material thermal limits still constrain practical engineering applications. The efficiency boundaries revealed by the Carnot cycle remain the gold standard for evaluating the performance of all thermodynamic systems to this day.

5.2. Optimization design of energy conversion systems and entropy production minimization strategies

Minimizing entropy production is essential for improving the thermodynamic efficiency of energy conversion systems. Irreversible losses such as friction, friction, temperature gradients, and chemical reactions—reduce energy quality, system lifespan, and increase emissions. Studies show that optimizing thermodynamic cycles, working fluids, and heat transfer paths can significantly reduce entropy production. For instance, advanced gas-steam combined cycles use multistage compression and expansion to limit pressure-related losses. In working fluid design, helium-liquid metal mixtures used in next-generation nuclear reactors lower entropy production to one-third of that in conventional systems, due to their low viscosity and high thermal conductivity. Bio-inspired fractal flow channel designs, modeled after leaf veins, further optimize the balance between heat transfer area and pressure losses.

In cogeneration systems, balancing heat use at high and low temperatures improves output and efficiency. Finite-time thermodynamics enables better optimization under variable conditions, such as in solar-ground source heat pump systems, where phase change control reduces entropy production by over 40%. In industrial waste heat recovery, countercurrent and cross-flow heat exchangers can reach 95% heat transfer efficiency, limiting entropy increase to under 5% of the total.Simulation studies suggest such optimizations can approach 80–85% of Carnot efficiency.

However, challenges remain. Multi-objective optimization must balance entropy reduction with cost and complexity. For example, recompression stages in supercritical CO_2 Brayton cycles improve efficiency by 4–6%, but increase costs by 15%. In wind-powered hydrogen production, fluctuating wind speeds create risks like compressor surge, making entropy control more difficult.

DOI: 10.54254/2753-8818/2025.GL24074

Looking ahead, AI and quantum computing are emerging as key tools. Deep reinforcement learning has reduced entropy production by 28% in fuel cell experiments by optimizing control strategies. Quantum computing enables faster, more complex entropy analysis, supporting multi-physics optimization tasks in minutes rather than weeks. These technologies are shifting the field from "entropy generation engineering" to "entropy generation intelligence," laying the foundation for ultra-efficient systems like the EU's "Horizon 2025" fusion project, which has already cut irreversible entropy production by 19%.

6. Conclusion

The second law of thermodynamics reveals the essential law of energy conversion in nature from micro and macro dimensions and provides a unified framework for understanding the irreversibility and energy efficiency limits of the system. At the micro level, the principle of entropy increase is explained by statistical mechanics: the Boltzmann entropy formula shows that the spontaneous evolution of an isolated system always favors macroscopic states with a larger number of molecular states (disorder). This microscopic randomness determines the irreversibility of macroscopic processes, such as the spontaneous transfer of heat from high temperature to low temperature, and the inability of fuel to spontaneously recover after combustion. On a macro level, the law guides engineering practice through entropy production analysis and becomes the core criterion for optimizing energy systems. Carnot cycle efficiency limit and entropy generation minimization strategies (such as heat exchanger topology optimization, waste heat step utilization) are aimed at reducing irreversible losses. For example, in gas turbine design, entropy production can be reduced by more than 20% by regulating combustion temperature and working fluid flow channels, significantly improving power generation efficiency. However, the real-time entropy control of dynamic systems, such as grid-connected renewable energy, still faces challenges, requiring multi-objective trade-offs combined with intelligent algorithms. The second law of thermodynamics not only defines the one-way arrow of time but also profoundly warns of the boundaries of energy use. From the perspective of sustainable development, only by following the constraint of entropy increase and approaching the reversible limit through technological innovation (such as near-zero entropy fuel cells and low-dissipation energy storage technology) can economic growth and ecological balance be coordinated. In the future, with the development of micro-scale entropy regulation techniques, such as molecular motor design, humans may open up new paths to greater energy efficiency in an irreversible universe.

References

[1] *Raffaele, P, Emilio, M.P, Abdelkader, A, Maxime, N. (2021). Conceptual polymorphism of entropy into the history: extensions of the second law of thermodynamics towards statistical physics and chemistry during nineteenth–twentieth centuries. Foundations of Chemistry, 23, 337-378.*

[2] *Sheldon, G. (2001). Boltzmann's Approach to Statistical Mechanics. Springer.com, Pis cataway, NJ, USA.*

[3] *Callen, H. B. (1985). Thermodynamics and an introduction to thermostatistics.Welly, New York.*

[4] *Prigogine, I. (1967). Introduction to Thermodynamics of Irreversible Processes. Interscience Publishers, New York.*

[5] *Schmid, A. (2020). The intersection of science and art in Viennese culture. Wiener Berichte über Naturwissenschaft in der Kunst, 15, 123–145.*

[6] *Boltzmann, L. (1877). Über die Beziehung zwischen dem zweiten Hauptsatze der mechanischen Wärmetheorie und der Wahrscheinlichkeitsrechnung. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Classe, 76, 373–435.*

[7] *Clausius, R. (1865). The Mechanical Theory of Heat. Kessinger Publishing, LLC, Kila, Montana.*

[8] *Kondepudi, D., Prigogine, I. (2015). Modern thermodynamics: From heat engines to dissipative structures (2nd ed.). Wiley.*

[9] *Clausius, R. (1865). On the application of the theorem of the equivalence of transformations to interior work. Annalen der Physik, 125(7), 353-400.*

DOI: 10.54254/2753-8818/2025.GL24074

[10] *Thomson, W. (1851). On the dynamical theory of heat. Transactions of the Royal Society of Edinburgh, 20(3), 261-298.*