Proceedings of ICBioMed 2025 Symposium: AI for Healthcare: Advanced Medical Data Analytics and Smart Rehabilitation
DOI: 10.54254/2753-8818/2025.AU24486

Optimization of Cell Culture Media Preparation: A Novel Cold-CO_2 Pre-equilibration Approach for Enhanced pH and Nutrient Stability

Bo Ma[1], Muttiah Barathan[1], Ping Yu[2], Min Hwei Ng[1], Jia Xian Law[1*]

[1]*Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia*
[2]*Center of Reproductive Medicine, Women's Hospital of Jiangnan University, Wuxi, China*
**Corresponding Author. Email: lawjx@ukm.edu.my*

Abstract. Maintaining optimal pH and nutrient stability in cell culture media is essential for ensuring cell viability and experimental reproducibility. Conventional methods, including direct warming (Method A) and pre-incubation in a CO_2 incubator (Method B), present limitations such as pH instability and nutrient degradation. This study introduces and evaluates a novel method (Method C), which involves equilibrating cell culture media overnight at 4°C in a 5% CO_2 environment prior to warming. Comparative analysis of these three methods indicated that Method C effectively maintains optimal pH levels while significantly preserving sensitive nutrients, including vitamin C and glutamine. Method C thus represents a promising alternative for optimizing cell culture conditions despite requiring additional preparation steps.

Keywords: cell culture media, pH, glutamine

1. Introduction

Cell culture media preparation is pivotal in experimental biology, directly impacting cellular growth, function, and reproducibility. Achieving optimal pH and nutrient integrity is critical, yet conventional preparation techniques often inadequately address these factors [1] . High temperatures reduce the solubility of CO_2. Under cultivation conditions at 37°C, a continuous supply of 5% CO_2 gas is required to maintain a constant dissolved CO_2 concentration, ensuring buffering capacity. However, prolonged cultivation still leads to an increase in pH [2,3]. At the same time, long-term cultivation at 37°C accelerates glucose consumption and lactic acid accumulation, as well as the breakdown of temperature-sensitive substances [4,5].

Method A, involving media warming at 37°C prior to use, results in elevated initial pH (~7.8), requiring lengthy equilibration periods. Method B, involving overnight incubation at 37°C with 5% CO_2, ensures immediate pH equilibrium but accelerates degradation of temperature-sensitive nutrients, notably Penicillin-Streptomycin and glutamine [6,7]. To mitigate these limitations, we propose Method C, a novel cold pre-equilibration method at 4°C under 5% CO_2, hypothesizing enhanced nutrient stability while maintaining optimal pH. This study evaluates the effectiveness of

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.AU24486

Method C in comparison to conventional methods regarding pH stability, nutrient integrity, and practical applicability.

2. Materials and methods

2.1. Cell culture media preparation

Media comprised DMEM supplemented with 10% FBS, 2 mM glutamine, and penicillin-streptomycin (100 U/mL).

2.2. Media pre-equilibration methods

Three distinct pre-equilibration protocols were systematically evaluated:

(1) Method A (Acute Warming):

Media was warmed at 37°C for 30 minutes immediately prior to use.

(2) Method B (Standard Overnight Incubation):

Media was incubated overnight (~16 hr) in a humidified 5% CO_2 incubator at 37°C.

(3) Method C (CO_2-Chilled Equilibration):

Overnight equilibration at 4°C with 5% CO_2(Figure 1), followed by 20-minute warming at 37°C prior to use.



Figure 1: Schematic diagram of the implementation process of Method C

2.3. pH measurement

pH was assessed immediately post-preparation, post-warming (Methods A and C), after 8-hour incubation (Method A), and prior to cell seeding (Methods B and C), utilizing a calibrated pH meter.

2.4. Nutrient stability

Penicillin-Streptomycin and glutamine concentrations were quantified via high-performance liquid chromatography (HPLC), assessing nutrient degradation across methods [8].

3. Results

3.1. pH stability

Method A demonstrated initial elevated pH (7.8), requiring ~8 hours for stabilization. Method B consistently provided optimal initial pH (7.4) but caused significant nutrient degradation. Method C

DOI: 10.54254/2753-8818/2025.AU24486

achieved stable optimal pH (7.4) prior to usage, comparable to Method B but with substantially improved nutrient stability.

3.2. Nutrient stability

Penicillin loss was significantly highest in Method B (~20%), intermediate in Method A (~10%), and lowest in Method C (~5%). Glutamine degradation mirrored these findings, with ~50% loss in Method B, whereas Method C effectively preserved approximately 90% of the original concentration.

3.3. Practical considerations

Method A offers simplicity and speed but requires extended post-seeding equilibration. Method B ensures immediate optimal pH but significantly compromises nutrient quality. Method C, despite additional setup requirements, offers optimal pH and superior nutrient preservation.

4. Discussion

This comparative analysis underscores the efficacy of Method C in balancing optimal media pH and nutrient stability. Its ability to significantly preserve penicillin and glutamine concentrations provides clear advantages, particularly for sensitive cellular models like primary and stem cells. Although Method C requires additional equipment and preparation, its benefits potentially outweigh these limitations. Future studies should extend this approach to varied cell types and media compositions to confirm broader applicability and long-term benefits.

5. Conclusion

Method C represents a novel and superior alternative to traditional cell culture media preparation, effectively maintaining optimal pH and nutrient stability, thus improving experimental consistency and cellular outcomes.

References

[1] van der Valk J, Brunner D, De Smet K, et al. Optimization of chemically defined cell culture media--replacing fetal bovine serum in mammalian in vitro methods. Toxicol In Vitro. 2010;24(4):1053-1063. doi:10. 1016/j. tiv. 2010. 03. 016

[2] Lee JT, Leng Y, Chow KL, et al. Cell culture medium as an alternative to conventional simulated body fluid. Acta Biomater. 2011;7(6):2615-2622. doi:10. 1016/j. actbio. 2011. 02. 034

[3] Wagener V, Virtanen S. Protective layer formation on magnesium in cell culture medium. Mater Sci Eng C Mater Biol Appl. 2016;63:341-351. doi:10. 1016/j. msec. 2016. 03. 003

[4] Yoshimura Y, Kikuiri T, Hasegawa T, et al. How much medium do you use for cell culture? Medium volume influences mineralization and osteoclastogenesis in vitro. Mol Med Rep. 2017;16(1):429-434. doi:10. 3892/mmr. 2017. 6611

[5] Koobotse MO, Schmidt D, Holly JMP, Perks CM. Glucose Concentration in Cell Culture Medium Influences the BRCA1-Mediated Regulation of the Lipogenic Action of IGF-I in Breast Cancer Cells. Int J Mol Sci. 2020;21(22):8674. Published 2020 Nov 17. doi:10. 3390/ijms21228674

[6] Hashizume T, Ozawa Y, Ying BW. Employing active learning in the optimization of culture medium for mammalian cells. NPJ Syst Biol Appl. 2023;9(1):20. Published 2023 May 30. doi:10. 1038/s41540-023-00284-7

[7] Flinck, M. , Kramer, S. H. , & Pedersen, S. F. (2018). Roles of pH in control of cell proliferation. Acta physiologica (Oxford, England), 223(3), e13068. https://doi. org/10. 1111/apha. 13068

DOI: 10.54254/2753-8818/2025.AU24486

[8] Vozniuk O, Kejík Z, Veselá K, et al. A Fast HPLC/UV Method for Determination of Ketoprofen in Cellular Media. ChemistryOpen. 2024;13(3):e202300147. doi:10. 1002/open. 202300147