Proceedings of ICBioMed 2025 Symposium: AI for Healthcare: Advanced Medical Data Analytics and Smart Rehabilitation
DOI: 10.54254/2753-8818/2025.AU25168

Research on Sleep Disorders and How Meditation Affects Sleep

Minghan Gong

Overseas Chinese Academy of Chiway Suzhou, Suzhou, China
inagong2008@icloud.com

Abstract. Sleep may appear to be the easiest thing for human beings; however, it is, in fact, a complex phenomenon influenced by many factors—including sleep disorders. Being unable to sleep normally brings great threat to the publication of human beings. In order to avoid many bad consequences, scientists have found ways to deal with them. This article discusses some different sleep disorders, ways to reduce their effects, and most importantly, the effects of mindfulness meditation on sleep. For individuals unfamiliar with this field, this study serves two primary purposes: to provide a general overview of existing knowledge and to help reduce societal misunderstandings about those who suffer from sleep disorders. Ultimately, this awareness may foster a more supportive and inclusive environment. This paper also addresses key questions such as: What are the types of sleep disorders? What is the relationship between meditation and sleep? Research suggests meditation can positively affect sleep, such as better sleep quality, easier to fall asleep, and reducing the negative effects of sleep disorders on people.

Keywords: Biology, Sleep disorders, Mindfulness Meditation on sleep, Brain Research

1. Introduction

Sleep disorders are illnesses that may cause you to be unable to fall asleep or keep you sleepy in the daytime. In this essay, the main subject covered is neurobiology, specifically sleep disorders, the method to deal with the disorders, and meditation as a big part of the research. Although scientists have discovered ways to relieve symptoms, some of the disorders —such as narcolepsy—remain incurable and can only be managed over time. For example, although patients with narcolepsy may experience symptom relief, there is still no definitive cure.

This paper serves as a summary and analysis of existing research findings, drawn from publicly available online resources. The primary research question centers on the impact of meditation on individuals with sleep disorders. The goal is not only to inform and educate readers but also to reduce stigma and prejudice surrounding those affected. Increased understanding may help foster empathy, reduce mental health burdens, and create a more inclusive society.

Awareness is especially important in everyday interactions. For instance, someone who suddenly falls asleep due to a sleep disorder might be misunderstood or discriminated against by those unaware of the condition. In extreme cases, such misunderstanding could even result in social

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.AU25168

exclusion or racism. Understanding the biological reasons behind such behavior can help promote equality and fairness.

The essay begins by defining the term "sleep disorders," which includes factors such as sleep quality (how restful sleep is), sleep timing (the ability to fall and stay asleep), and sleep quantity (how much sleep a person gets). Importantly, a single sleepless night does not indicate a disorder. Sleep disorders are suspected only when irregularities occur frequently, cause fatigue even after sufficient sleep duration, and interfere with daily functioning [1]. Additionally, the lack of public awareness and diagnostic clarity often delays professional intervention, worsening the condition over time.

2. Overview of sleep disorders

By understanding the most common sleep disorders and their impact on individuals, readers may begin to reduce the bias or prejudice they might have previously held toward those affected. The goal is not only to promote a more accepting society but to contribute to a world where differences are respected and understood. Following this overview, the paper will also discuss treatment options for sleep disorders to enhance readers' knowledge and equip them—and potentially their friends or family members—with accessible remedies and support strategies.

2.1. Insomnia

Insomnia is arguably the most prevalent sleep disorder among the global population. According to the World Health Organization, the percentage of people worldwide who have difficulties falling asleep is 27%, less than half of the people, but still significant data [2]. This widespread issue highlights the importance of early detection and intervention.

Insomnia is characterized by difficulty falling asleep or staying asleep despite having adequate opportunity and desire to sleep. The condition is largely beyond voluntary control and can significantly affect daily life. For instance, an individual who cannot fall asleep the night before an important event may find their entire next day compromised by fatigue and impaired focus. Consequently, insomnia is often accompanied by excessive daytime sleepiness, reduced cognitive function, and emotional instability. It is more common among older adults.

Understanding the causes of insomnia is crucial. Common contributing factors include stress, irregular sleep schedules, poor sleep hygiene, mental health conditions (e.g., anxiety and depression), physical pain or illness, certain medications, neurological issues, and other sleep disorders. Often, insomnia results from a combination of these elements.

When symptoms occur at least three times per week over a period of three months, the condition can be diagnosed as chronic insomnia, one of the most common subtypes [3]. Early recognition and intervention are essential in preventing long-term negative outcomes.

2.2. Sleep apnea

Sleep apnea is a more severe sleep disorder than insomnia and can pose serious health risks. It is classified as a breathing-related disorder that disrupts normal respiration during sleep. Individuals with sleep apnea often experience loud snoring, gasping, or choking during the night, leading to frequent awakenings.

The consequences are profound: fragmented sleep results in daytime fatigue, reduced concentration, and an increased risk of cardiovascular and mental health problems. People with

DOI: 10.54254/2753-8818/2025.AU25168

untreated sleep apnea may also suffer from mood disorders due to chronic exhaustion and disturbed sleep.

There are two main types of sleep apnea, both sharing similar symptoms but differing in origin:

Obstructive Sleep Apnea (OSA): This type occurs when the soft tissues in the throat and mouth relax excessively during sleep, causing a blockage of the upper airway and momentary interruptions in breathing.

Central Sleep Apnea (CSA): This less common form is neurological in nature. It occurs when the brain fails to send proper signals to the muscles that control breathing, leading to breathing pauses.

In both cases, individuals may experience morning headaches, excessive daytime sleepiness, and irritability. Despite its seriousness, sleep apnea is highly underdiagnosed, with many sufferers unaware of the condition until it is observed by someone else or leads to secondary health problems.

Narcolepsy does not typically disrupt nighttime sleep but can have severe effects during the daytime, particularly in critical or high-stakes situations. It can interfere with work, study, or social interactions—such as during meetings or client presentations—creating significant challenges in everyday life.

The primary symptom of narcolepsy is excessive daytime sleepiness (EDS), even after a full night's rest. In more severe cases, individuals may experience "sleep attacks," which are sudden, uncontrollable episodes of falling asleep that last for a few minutes. These episodes can occur at any time, regardless of whether the person is standing, working, exercising, or simply sitting. Because of their unpredictability, sleep attacks can disrupt important tasks and pose serious risks in situations that require attention and physical coordination.

This disorder is especially dangerous when individuals are engaged in activities such as cooking, climbing, operating machinery, or driving. For example, falling asleep unexpectedly while a stove is on could lead to fires or injury. As a result, people with narcolepsy must take precautions to avoid engaging in high-risk tasks without appropriate supervision or safety measures.

Narcolepsy may also co-occur with cataplexy, a condition characterized by a sudden loss of muscle tone, often triggered by strong emotions such as laughter or surprise. During cataplexy episodes, individuals may collapse or slump over as their body temporarily loses control, which is both physically dangerous and socially distressing.

Narcolepsy is caused by disruptions in the brain's ability to regulate the sleep-wake cycle, often linked to a deficiency in the neurotransmitter hypocretin (also known as orexin), which plays a key role in maintaining wakefulness and preventing REM sleep from occurring at inappropriate times.

In summary, insomnia, sleep apnea, and narcolepsy are three of the most common sleep disorders, each presenting unique challenges to affected individuals. Other notable sleep-related conditions include restless legs syndrome (RLS), parasomnias (such as sleepwalking or night terrors), hypersomnia (excessive sleepiness), shift work disorder, and non-24-hour sleep-wake disorder. While the symptoms and causes differ, all of these disorders significantly impact a person's quality of life and daily functioning.

3. Existing remedies for sleep disorders

There are a variety of existing methods available to help manage sleep disorders. While some conditions, such as narcolepsy, currently have no complete cure, their symptoms can often be alleviated through treatment. Many disorders share general treatment strategies, such as medications and therapy, though the specific forms may vary.

For a common disorder like insomnia, treatments include cognitive behavioral therapy for insomnia (CBT-I), prescription or over-the-counter sleep aids, and lifestyle adjustments. Sleep aids

DOI: 10.54254/2753-8818/2025.AU25168

are a category of medications that help individuals fall asleep by inducing drowsiness through their sedative effects. While effective in the short term, they must be used cautiously due to the risk of dependence or side effects.

For sleep apnea, a widely used and effective treatment is CPAP (Continuous Positive Airway Pressure) therapy. This involves a machine that delivers a steady stream of air pressure through a mask, keeping the upper airway open during sleep. By ensuring continuous airflow into the lungs, CPAP therapy helps prevent breathing interruptions, enhances sleep quality, and significantly reduces health risks associated with sleep apnea, such as heart attack and stroke [4].

Other disorders have more specific treatments. For instance, Restless Legs Syndrome (RLS) can often be managed through dietary adjustments, iron supplements, or certain medications that address neurological causes. In the case of parasomnias—disorders that involve unusual behaviors during sleep, such as sleepwalking or night terrors—safety becomes the top priority. It is essential to create a secure environment to prevent injury to the person or their bed partner during episodes.

Excessive daytime sleepiness is not a disorder in itself but is a symptom associated with various sleep disorders. Because its underlying causes vary between individuals, diagnostic tests and individualized sleep studies conducted by medical professionals are essential for effective treatment.

Shift Work Disorder and Non-24-Hour Sleep-Wake Disorder both involve irregular sleep patterns, though in different ways. In Shift Work Disorder, symptoms result from disruptions caused by non-traditional work hours. The key solution is to maximize sleep quality during off-hours, including the use of sleep hygiene techniques or light therapy. In contrast, Non-24-Hour Sleep-Wake Disorder is often seen in blind individuals and is caused by a misalignment between the body's internal clock and the 24-hour day. Treatment typically involves behavioral therapies and melatonin supplementation. Melatonin, a hormone that regulates the sleep-wake cycle, can be taken at the same time each night to help the body synchronize with a regular schedule [5].

Although these treatments are helpful, ongoing research and technological advancements continue to improve outcomes. Future innovations may offer more effective and accessible solutions, allowing a greater number of people to regain control over their sleep and overall well-being.

Having reviewed standard medical and therapeutic interventions, the next section will focus on meditation, a particularly intriguing and non-invasive technique that has shown promise in supporting individuals with sleep disorders.

4. The effects of mindfulness meditation on sleep

4.1. Definitions of meditation

To understand how meditation influences sleep, it is essential to first define what mindfulness meditation entails. Meditation is a research-backed technique for reducing stress and promoting mental and physical well-being by inducing measurable changes in the brain and body. It can be defined as the practice of training one's attention to achieve a mental state characterized by calm concentration and positive emotions.

Mindfulness meditation, one of the most widely practiced forms, consists of two main components: attention and acceptance. The attention aspect involves deliberately focusing on the present moment—this may include awareness of one's breath, thoughts, physical sensations, and emotions. The acceptance aspect requires individuals to observe these experiences without judgment. Rather than reacting emotionally or intellectually, the individual simply acknowledges the thoughts and sensations and then lets them pass [6].

DOI: 10.54254/2753-8818/2025.AU25168

These mindfulness-based interventions (MBIs)—including Mindfulness-Based Stress Reduction (MBSR) and Mindfulness-Based Cognitive Therapy (MBCT)—have been shown to reduce pain, fatigue, and stress in individuals suffering from chronic pain, which in turn can alleviate related sleep disorders [7].

4.2. The physiological and psychological mechanisms of meditation

The effects of mindfulness meditation on sleep have been widely studied, and results consistently demonstrate its positive impact. In several research programs involving individuals with sleep disturbances caused by different disorders, participants who practiced mindfulness meditation experienced significant improvements in sleep quality compared to those in inactive control groups.

Although these findings are promising, the precise mechanisms through which mindfulness meditation improves sleep are still being explored. One widely accepted hypothesis suggests that these benefits are closely related to the brain, which governs both emotional responses and physiological states. The following discussion is based on insights gathered from scholarly articles and scientific resources.

At its core, meditation involves turning inward to observe one's thoughts and emotions, a seemingly simple act that produces substantial benefits. One key explanation is that mindfulness meditation reduces the body's stress response, which plays a critical role in many sleep disorders. According to researchers, mindfulness practice can "dial down" the stress response, leading to improved bodily function. Chronic stress—a common consequence and contributor to sleep disorders—can impair immune function and exacerbate various health conditions. By reducing stress, mindfulness meditation may thus produce wide-ranging positive effects throughout the body [7].

But how exactly does it reduce stress? Psychological studies suggest that meditation affects two primary stress pathways in the brain. Specifically, mindfulness alters brain structure and neural activity in regions associated with attention regulation and emotional control. These neurological changes have been linked to reduced negative thinking, emotional reactivity, and anxiety. Moreover, individuals who meditate tend to focus more on the present moment rather than ruminating on the past or worrying about the future, which leads to a significant decrease in stress levels [8].

4.3. Common meditation practices

There are nine common types of meditation practices that individuals may adopt. These include mindfulness meditation—which is the primary focus of this essay—spiritual meditation, focused meditation, movement meditation, mantra meditation, transcendental meditation, progressive relaxation, loving-kindness meditation, and visualization meditation. Each form offers different techniques and goals, but all share the common purpose of enhancing mental clarity, reducing stress, and promoting emotional well-being.

4.4. Explanation

This leads to an important question: what are the two distinct stress response pathways, and how does meditation influence brain structure and activity in specific regions?

One of these pathways begins in the hypothalamus, a small but vital structure in the brain that regulates many bodily functions. When a stressor is perceived, the hypothalamus sends a signal to the pituitary gland, which then communicates with the adrenal glands located on top of the kidneys

DOI: 10.54254/2753-8818/2025.AU25168

[8]. The adrenal glands produce a number of essential hormones, including cortisol, aldosterone, and adrenaline, which are responsible for regulating metabolism, blood pressure, and the body's stress response.

Cortisol, in particular, plays a central role in managing stress. It helps regulate blood pressure, raises blood sugar levels, and contributes to controlling the body's sleep-wake cycle [9]. Therefore, it is hypothesized that mindfulness meditation may help regulate cortisol levels, thereby aiding in stress reduction and improving sleep quality.

Further research has been conducted to better understand the role of cortisol in these physiological processes, especially in relation to sleep regulation and the stress response system.

4.5. Overview

IIn summary, the adrenal glands operate under the control of the hypothalamus and the pituitary gland. The hypothalamus participates in hormonal regulation by producing corticotropin-releasing hormone (CRH) and antidiuretic hormone (ADH), also known as vasopressin. These hormones stimulate the pituitary gland to release adrenocorticotropic hormone (ACTH), which in turn prompts the adrenal glands to produce corticosteroids such as cortisol and aldosterone [9].

As a result, when individuals engage in mindfulness meditation—paying attention and practicing acceptance of their internal states—it may activate the hypothalamus to initiate this hormonal cascade. This ultimately reduces stress and contributes to better regulation of the sleep-wake cycle.

Additionally, the kidneys contribute to the regulation of aldosterone, and the sympathetic nervous system influences the release of adrenaline and noradrenaline from the adrenal glands. Together, these systems interact with meditative practices to support stress regulation and sleep management [10].

5. Conclusion

To conclude, sleep disorders are varied in form, but they all involve difficulties in falling asleep, staying asleep, or achieving restful sleep. It is evident that treatments differ depending on the causes of the disorder, and each approach requires deeper investigation for more effective outcomes. Among the various treatment options, meditation has emerged as a widely used and accessible strategy, particularly for certain types of sleep disturbances. This is not only due to its convenience —requiring no medication or special equipment—but also because individuals can practice it anywhere, provided they can maintain focus.

This essay has explored the physiological mechanisms behind mindfulness meditation as a treatment for sleep disorders. In summary, mindfulness meditation stimulates the hypothalamus to produce hormones, which in turn signal the adrenal glands to release stress-regulating hormones such as cortisol. Additionally, mindfulness meditation engages the sympathetic nervous system, contributing further to stress reduction. These processes together can improve sleep quality.

Furthermore, the psychological benefits of reduced stress and increased emotional regulation may encourage trust and support from patients' families or friends, creating an environment that facilitates continued meditation practice and reinforces its positive effects.

However, this study is limited by the number of sources consulted, and future research should incorporate a broader range of scientific literature and empirical studies to deepen the understanding of mindfulness meditation's role in sleep health.

DOI: 10.54254/2753-8818/2025.AU25168

References

[1] Chavda, V., Chaurasia, B., Umana, G. E., Tomasi, S. O., Lu, B., & Montemurro, N. (2022). Narcolepsy—A neuropathological obscure sleep Disorder: A Narrative review of Current literature. Brain Sciences, 12(11), 1473. https: //doi.org/10.3390/brainsci12111473
[2] World Health Organization: WHO. (2020, January 27). Disability. https: //www.who.int/health-topics/disability
[3] McNamara, S., Spurling, B. C., & Bollu, P. C. (2025, March 28). Chronic insomnia. StatPearls - NCBI Bookshelf. https: //www.ncbi.nlm.nih.gov/books/NBK526136/
[4] Professional, C. C. M. (2024, July 12). CPAP machine. Cleveland Clinic. https: //my.clevelandclinic.org/health/treatments/22043-cpap-machine
[5] Casey O. Diekman , Amitabha Bose. (2022). Beyond the limits of circadian entrainment: Non-24-h sleep-wake disorder, shift work, and social jet lag. ScienceDirect. https: //www.sciencedirect.com/science/article/abs/pii/S0022519322001461
[6] Rusch, H. L., Rosario, M., Levison, L. M., Olivera, A., Livingston, W. S., Wu, T., & Gill, J. M. (2018). The effect of mindfulness meditation on sleep quality: a systematic review and meta-analysis of randomized controlled trials. Annals of the New York Academy of Sciences, 1445(1), 5–16. https: //doi.org/10.1111/nyas.13996
[7] Mindfulness meditation. (2019, October 30). American Psychological Association. Retrieved March 10, 2025, from https: //www.apa.org/topics/mindfulness/meditation
[8] Smith, S. M., & Vale, W. W. (2006). The role of the hypothalamic-pituitary-adrenal axis in neuroendocrine responses to stress. Dialogues in Clinical Neuroscience, 8(4), 383–395. https: //doi.org/10.31887/dcns.2006.8.4/ssmith
[9] Professional, C. C. M. (2025, February 10). Adrenal Gland. Cleveland Clinic. https: //my.clevelandclinic.org/health/body/23005-adrenal-gland
[10] Breus, M. J., PhD. (2020b, April 10). How poor sleep can affect cortisol levels. Psychology Today. https: //www.psychologytoday.com/us/blog/sleep-newzzz/202004/the-effects-cortisol-your-sleep