Proceedings of ICBioMed 2025 Symposium: Computational Modelling and Simulation for Biology and Medicine
DOI: 10.54254/2753-8818/2025.LD25430

Adaptation of Fish to the Extreme Environment in the Deep Sea

Luyao Mao

School of Life Sciences, Lanzhou University, Lanzhou, China
320220927410@lzu.edu.cn

Abstract. Deep-sea adaptation research has evolved from initial faunal descriptions to decoding the mechanistic basis of extremophile survival. Genomic analyses reveal conserved adaptive strategies, yet critical gaps persist, particularly regarding the hadal zone's extreme pressures, temperatures, and darkness. This paper examines the synergistic molecular, sensory, and evolutionary adaptations enabling teleost fish, primarily hadal liparids, to withstand this abiotic triad: it specifically investigates macromolecular stabilization under concurrent thermal and baric stress, conserved patterns in sensory compensation across depth gradients, and the contributions of exaptation versus de novo evolution. The analysis employs a multidisciplinary framework integrating comparative genomic data from 27 deep-sea teleost species with in vitro validation of pressure-temperature responsive proteins. This is augmented by phylogenetically informed trait evolution analysis and metabolic modeling under nutrient-limited conditions to establish cross-validated mechanistic insights. The synthesis reveals key innovations in protein conformations and sensory specialization, identifies patterns of conserved evolutionary pathways, and highlights the role of both exaptation and novel traits. Understanding these adaptations advances theoretical models of biological tolerance limits, which also offers transformative applications for biomimetic enzyme stabilization, deep-sea conservation vulnerability assessments, and frameworks for extraterrestrial life detection in subglacial oceans.

Keywords: Hadal adaptation, Pressure-temperature stress, Molecular mechanisms, Comparative genomics

1. Introduction

Deep-sea adaptation research has progressed from initial faunal documentation to mechanistic decoding of extremophile survival. Genomic analyses now reveal conserved strategies across taxa, including pressure-resistant protein conformations and sensory specialization. Despite advances, critical gaps persist regarding in situ validation of adaptation models, interactions between concurrent stressors, and evolutionary trajectories of non-model organisms. The hadal zone (6,000–11,000 m), representing the planet's most extreme marine environment, remains particularly underexplored mechanistically.

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.LD25430

This review examines the synergistic adaptation mechanisms enabling teleost fish—primarily hadal-endemic liparids—to withstand the abiotic triad defining deep-sea habitats: hydrostatic pressure (≤1,000 atm), near-freezing temperatures (0–4°C), and perpetual darkness.

We further advocate standardizing hadal simulation methodologies to enable cross-study comparisons and accelerate discovery of biomimetic solutions.

This paper specifically investigates molecular innovations that stabilize macromolecular structures under concurrent thermal and baric stress, determining whether sensory compensation mechanisms follow conserved evolutionary patterns along depth gradients, and evaluating the relative contributions of exaptation versus de novo evolution in deep-sea adaptations. The analytical framework synthesizes comparative genomic data from 27 deep-sea teleost species with in vitro validation of pressure-temperature responsive proteins, while incorporating phylogenetically informed trait evolution analysis and metabolic modeling under nutrient-limited conditions to establish cross-validated mechanistic insights. Understanding these adaptations advances theoretical models of biological tolerance limits, with transformative applications including piezolyte-inspired enzyme stabilization reagents for biotechnology, vulnerability assessments for deep-sea mining conservation strategies, and extraterrestrial life detection frameworks relevant to subglacial ocean astrobiology.

2. Low-temperature adaptation mechanisms

The constant, frigid environment of the deep sea has driven the evolution of sophisticated cold-adaptation strategies. A fundamental challenge is maintaining membrane fluidity, as cold temperatures induce membrane rigidification ("cold-hardening"). Deep-sea fishes counteract this through compositional changes in their cellular membranes. Notably, the amplification of genes like acaa1 significantly elevates the proportion of polyunsaturated fatty acids (PUFAs) within membrane phospholipids [1-3]. The kinks introduced by PUFA double bonds prevent tight packing of lipid molecules, thereby preserving essential membrane fluidity and functionality even at near-freezing temperatures. Crucially, this adaptation also serendipitously mitigates the membrane-rigidifying effects of high hydrostatic pressure, demonstrating a synergistic solution to dual stressors.

Protein function undergoes significant adaptive evolution for low-temperature operation. Antarctic icefish (family Channichthyidae), inhabiting the coldest waters, exemplify this. Mutations at key residues (S160A and V259M) within their visual pigment, rhodopsin, significantly lower the activation energy barrier required for the protein's conformational shift upon photon absorption [4]. This optimization allows efficient phototransduction necessary for vision despite the drastically slowed molecular kinetics inherent to near-ice-point temperatures. Beyond specific proteins, fundamental cellular processes like transcription are also fine-tuned. Deep-sea fishes, including those from the hadal zone, exhibit convergent mutations in the rtf1 gene [3]. This gene encodes a factor crucial for transcription elongation, and these mutations appear to optimize this process under the constraints of low temperature, ensuring adequate gene expression.

Energy conservation is paramount in the cold, nutrient-poor deep sea. Fishes have evolved strategies at multiple levels. Neurologically, there is evidence for structural simplification within neural networks, reducing the energetic costs associated with maintaining complex brain function where high cognitive processing may be less critical [3,5]. Genetically, deep-sea fishes often exhibit lower genomic mutation rates compared to shallow-water relatives [3,5]. This reduction potentially minimizes the energetic burden of DNA repair mechanisms and the production of deleterious mutations, representing a metabolic saving strategy over evolutionary timescales. Collectively, these

DOI: 10.54254/2753-8818/2025.LD25430

findings illuminate the profound molecular underpinnings of cold adaptation and reveal striking evolutionary parallels between deep-sea and polar organisms facing similar thermal challenges.

3. High-pressure adaptation mechanisms

Adapting to hydrostatic pressures that can exceed 1000 times that at the sea surface (1 kbar or 100 MPa) is arguably the most unique challenge for hadal vertebrates. Research understanding has shifted considerably. Early theories emphasized the critical role of the osmolyte trimethylamine N-oxide (TMAO) in counteracting pressure-induced protein denaturation. TMAO stabilizes protein structure by preferential exclusion from the protein surface, effectively promoting a compact, folded state. However, a critical limitation emerged, that TMAO's beneficial effects are concentration-dependent, but its accumulation simultaneously increases intracellular osmotic pressure. This osmotic constraint appears to set a physiological depth limit of approximately 8200 meters for teleost fish distribution, beyond which the required TMAO concentration becomes osmotically untenable [6].

This realization spurred the search for alternative pressure-adaptation mechanisms, revealing a complex, multi-gene "defense" system. A landmark discovery involved the hadal snailfish (Pseudoliparis swirei). Genomic analysis revealed tandem duplication of the ftlh27 (ferritin, heavy polypeptide-like 27) gene [7]. Functional experiments demonstrated that this duplication confers a significantly enhanced capacity (up to 40% reduction) to mitigate reactive oxygen species (ROS) damage induced by high pressure [7]. Pressure can exacerbate oxidative stress, making this antioxidant defense crucial. Furthermore, genes involved in maintaining cellular architecture show strong signatures of positive selection. Genes encoding vinculin (vcl) and phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit (pik3), key players in cytoskeletal organization and signal transduction related to cell shape and adhesion, exhibit adaptations specifically beneficial for preserving cellular integrity and morphology under extreme compression [2]. This prevents pressure-induced cytoskeletal collapse and membrane damage.

Buoyancy regulation under high pressure presents another critical challenge. A long-standing hypothesis proposed that deep-sea fishes progressively reduce bone density with increasing depth to achieve neutral buoyancy. However, sophisticated micro-computed tomography (micro-CT) imaging combined with rigorous phylogenetic comparative analysis has overturned this view [8]. Studies on diverse groups like grenadiers (Macrouridae) demonstrate that successful deep-sea species, including those at hadal depths, often achieve neutral buoyancy through specialized swim bladders. These bladders are filled with low-density gases maintained under pressure, but crucially, some species, like certain macrourids, retain functional swim bladders lined with highly reflective guanine crystals, even at great depths [8]. This mechanism provides efficient buoyancy without necessitating wholesale skeletal reduction. These combined discoveries establish a contemporary model of high-pressure adaptation reliant on synergistic interactions between antioxidant systems, cytoskeletal stabilizers, and specialized buoyancy organs, all underpinned by specific genetic modifications.

4. Darkness adaptation mechanisms

The perpetual darkness of the deep sea below the photic zone has driven profound evolutionary changes in sensory systems, particularly vision, and necessitated compensatory enhancements in other senses. Visual adaptation is multifaceted and remarkably sophisticated. Contrary to the traditional assumption that deep-sea fishes possess monochromatic vision, spectral tuning of visual pigments allows for discrimination within the limited light available. The silver spinyfin (Diretmus

DOI: 10.54254/2753-8818/2025.LD25430

argenteus) possesses an extraordinary repertoire of 38 opsin genes, with at least 14 distinct rod opsin genes expressed in its retina [9]. This genetic expansion allows for the expression of multiple rhodopsin photopigments tuned to different wavelengths. Crucially, these pigments cover two primary light sources: the blue-green bioluminescence produced by many deep-sea organisms (peaking between 420nm and, 520 nm) and the extremely dim residual downwelling sunlight, filtered to a narrow blue band (around 432-507 nm) [10]. This opsin diversity provides the potential for dichromatic or even polychromatic vision in near-total darkness. Structural adaptations further maximize photon capture. Deep-sea fish retinas often feature a banked (multilayered) rod structure, dramatically increasing the photoreceptor density per unit area. This is coupled with an exceptionally high density of rhodopsin molecules within the rod outer segments, that the densities can reach twice that found in shallow-water fish [5]. Finally, a reflective tapetum lucidum layer behind the retina acts like a mirror, reflecting photons back through the photoreceptor layer for a second chance at absorption, significantly enhancing sensitivity.

Compensatory sensory augmentation is vital in the absence of light. Hearing becomes a critical long-range sense. The hadal snailfish (P. swirei) shows significantly elevated expression of the tmc1 (transmembrane channel-like 1) gene in its inner ear [7]. This gene is essential for mechanotransduction in auditory hair cells, suggesting enhanced auditory sensitivity for detecting sounds like water movements or prey-generated noises. Olfaction also undergoes enhancement. Deep-sea cusk-eels (Ophidiidae) exhibit substantial expansions in their repertoire of olfactory receptor genes [3]. This expansion likely allows for the detection of a broader spectrum of chemical cues dissolved in the water, essential for locating food, mates, or navigating in the dark. These enhancements in auditory and olfactory capabilities represent a clear case of cross-modal sensory compensation [3].

The perpetual darkness also fundamentally alters circadian biology. Deep-sea fishes retain the core molecular machinery of the circadian clock (genes like clock, bmal, cry and per), indicating the importance of maintaining an internal rhythm for coordinating physiology and behavior [7]. However, key light-responsive elements and pathways that normally synchronize (entrain) this internal clock to the external day/night cycle in shallow waters are often degraded or lost [7]. This results in a functional decoupling: the endogenous circadian oscillator continues to run, but its phase is no longer reset by environmental light cues. The timing of this internal rhythm may instead be influenced by other cyclic factors like tidal currents or periodic vertical migrations of prey.

5. Interconnected mechanisms and research horizons

The adaptations to low temperature, high pressure, and darkness are not isolated but exhibit significant synergy and shared molecular strategies [1-3]. Similarly, the remarkable diversity of rod opsins enhances visual sensitivity in darkness and compensates for the reduced kinetics of visual transduction proteins (like rhodopsin) caused by low temperatures [4,10]. Gene duplication emerges as a fundamental evolutionary tactic across environmental challenges. Examples include the ftlh27 tandem duplication for antioxidant defense under pressure [7], and potentially duplications in genes like cldnj (involved in cell junction integrity) contributing to adaptation across multiple stressors [7], highlighting its role as a versatile engine for evolutionary innovation in the deep sea.

Research methodologies have become increasingly integrative, driving recent progress. Biochemical quantification, such as precise measurement of TMAO concentrations across depth gradients [6], provided foundational insights. Advanced gene editing techniques (e.g., CRISPR-Cas9) now allow direct functional validation of candidate genes, as demonstrated for ftlh27 in

DOI: 10.54254/2753-8818/2025.LD25430

pressure adaptation [3,7]. Detailed microscopic dissection and imaging techniques, including micro-CT, have revolutionized our understanding of anatomical adaptations like swim bladder structure and retinal organization [5,8]. Computational approaches, particularly molecular dynamics simulations, offer powerful tools to model how specific mutations (like those in rhodopsin or cytoskeletal proteins) alter protein structure, dynamics, and function under simulated high pressure or low temperature [10].

6. Conclusion

Despite significant advances, critical limitations and exciting frontiers remain. A major gap is the scarcity of in situ physiological and biochemical data collected under genuine deep-sea conditions; most experiments are necessarily conducted on retrieved specimens under surface pressure and temperature. Key future research directions are therefore multifaceted and interconnected: Firstly, there is an urgent need to systematically investigate pressure-temperature interactions by probing the combined effects of high pressure and low temperature on fundamental processes like protein folding stability, enzyme kinetics, and membrane phase behavior, specifically elucidating how these synergistic stressors interact at the molecular level. Complementing this molecular focus. Secondly, research must prioritize Behavioral Ecology in Darkness to validate the ecological significance of sensory adaptations inferred from genomics and anatomy. This requires answering critical questions about how deep-sea fishes actually utilize their enhanced senses, such as spectral discrimination in bioluminescence and enhanced olfaction or audition, for essential behaviors like foraging, communication, predator avoidance, and navigation in their natural habitat; deploying unobtrusive deep-sea observation platforms is crucial for this endeavor. Thirdly, a deeper understanding of Energetics and Metabolism is essential, demanding the quantification of the energetic costs inherent to high-pressure life and the efficiency of deep-sea metabolic adaptations, potentially achievable through the development of advanced respirometry techniques adapted for pressure. Finally, as an emerging frontier, exploring Hadal Microbial Symbioses will illuminate the potential role of symbiotic microorganisms within the gut or specialized organs, not only for nutrient acquisition in the oligotrophic deep but also potentially for detoxification or stress mitigation mechanisms at extreme depths. The molecular adaptations of deep-sea fishes provide profound insights into the limits of vertebrate life and the ingenious solutions evolution can craft. Studying these extremophiles not only satisfies fundamental biological curiosity but also offers valuable paradigms for understanding stress resilience, potentially informing fields from biotechnology to medicine. Their survival strategies, forged in one of Earth's most hostile environments, stand as a testament to the power of natural selection.

References

[1] Wang, K., et al. (2019). Morphology and genome of a snailfish from the Mariana Trench provide insights into deep-sea adaptation. Nat Ecol Evol, 3(5): 823-833.
[2] Bo, J., et al. (2022). Molecular Mechanisms of the Convergent Adaptation of Bathypelagic and Abyssopelagic Fishes. Genome Biol Evol, 14(8).
[3] Xu, H., et al. (2025). Evolution and genetic adaptation of fishes to the deep sea. Cell 188(5): 1393-1408 e1313.
[4] Castiglione, G. M., et al. (2023). Adaptation of Antarctic Icefish Vision to Extreme Environments. Mol Biol Evol, 40(4).
[5] Wagner, H. J., et al. (1998). The eyes of deep-sea fish. II. Functional morphology of the retina. Prog Retin Eye Res, 17(4): 637-685.
[6] Yancey, P. H., et al. (2014). Marine fish may be biochemically constrained from inhabiting the deepest ocean depths. Proc Natl Acad Sci U. S. A., 111(12): 4461-4465.

DOI: 10.54254/2753-8818/2025.LD25430

[7] Xu, W. J., et al. (2023). Chromosome-level genome assembly of hadal snailfish reveals mechanisms of deep-sea adaptation in vertebrates. Elife, 12.
[8] Martin, R. P., et al. (2022). Bone Density Variation in Rattails (Macrouridae, Gadiformes): Buoyancy, Depth, Body Size, and Feeding. Integr Org Biol, 4(1): obac044.
[9] Warrant, E. J. and N. A. Locket (2004). Vision in the deep sea. Biol Rev Camb Philos Soc, 79(3): 671-712.
[10] Musilova, Z., et al. (2019). Vision using multiple distinct rod opsins in deep-sea fishes. Science, 364(6440): 588-592.