Proceedings of ICBioMed 2025 Symposium: AI for Healthcare: Advanced Medical Data Analytics and Smart Rehabilitation
DOI: 10.54254/2753-8818/2025.AU27668

Stem Cell Rejuvenation via Heterochronic Parabiosis: A Literature Review and Case Analysis

Hanyu Dou

Changchun Shiyi High School, Changchun, China
3988098851@qq.com

Abstract. Aging is a complex biological process marked by progressive tissue function decline and reduced regenerative capacity, with stem cell exhaustion being a key driver. Heterochronic parabiosis (HP), which surgically connects the circulatory systems of young and old organisms, has become a vital tool for exploring systemic rejuvenation, especially its role in reversing age-related stem cell dysfunction via young blood factors. This review synthesizes current knowledge on HP, analyzes key studies (including one using single-cell transcriptomics to demonstrate multi-tissue rejuvenation), and addresses controversies around factors like GDF11 and TIMP2. It also discusses challenges such as species translatability and ethical concerns, and concludes by proposing future directions like refined factor-based therapies and combinatorial approaches targeting stem cell niches.

Keywords: Stem cell rejuvenation, Heterochronic parabiosis, Aging

1. Introduction

Aging manifests through hallmarks like genomic instability and altered intercellular communication, with stem cell depletion being a central pillar. The heterochronic parabiosis (HP) model, pioneered by Conboy et al., revealed that young systemic environments can revitalize aged tissues, sparking interest in "rejuvenation factors"—a discovery that has since driven extensive research into leveraging systemic cues for anti-aging interventions. As a vital tool for exploring systemic rejuvenation, HP has provided critical insights into how young blood factors may reverse age-related stem cell dysfunction, with studies even using single-cell transcriptomics to demonstrate multi-tissue rejuvenation effects. However, the field also faces ongoing controversies, such as debates around the roles of specific factors like GDF11 and TIMP2.

This chapter (and the broader review) aims to evaluate HP's scientific impact, critique key studies in the field, discuss barriers to clinical translation (including species translatability and ethical concerns), and lay the groundwork for exploring future research directions. By synthesizing current knowledge on HP and its relevance to stem cell aging and systemic rejuvenation, this paper seeks to clarify the model's value in advancing anti-aging research while highlighting the challenges that must be addressed to move toward practical applications like factor-based therapies or stem cell niche-targeted interventions.

© 2026 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.AU27668

2. Literature review

2.1. Mechanisms of stem cell aging

Aging stem cells undergo progressive functional decline due to cumulative intrinsic damage. DNA damage accumulation leads to increased double-strand breaks and compromised repair mechanisms, triggering p53-mediated quiescence or apoptosis that depletes regenerative reserves [1]. Telomere attrition in hematopoietic and muscle satellite cells induces replicative senescence, with telomerase deficiency causing premature aging in model systems. Concurrently, epigenetic alterations, including DNA methylation loss at polycomb targets and reduced H3K27me3 marks, disrupt stem cell identity, though cellular reprogramming demonstrates these changes are potentially reversible. Together, these intrinsic impairments create a molecular signature of stem cell aging characterized by genomic instability, proliferative arrest, and identity loss.

The aged microenvironment severely impairs the functions of stem cells through multiple mechanisms. One of the key factors is the "senescence-associated secretory phenotype (SASP)," where senescent cells release pro-inflammatory cytokines (such as IL-6 and TNF-α), thereby interfering with the activities of adjacent stem cells - the use of senescence cell clearance agents (such as dasatinib + vincristine) to remove these cells has been proven to enable the rejuvenation of aged tissues. Additionally, disrupted signaling pathways also play a crucial role: in aged intestinal stem cells (ISCs), the over-activated "Wnt/β-catenin" signaling pathway promotes differentiation into secretory cell lineages, thereby reducing the stem cell pool; in neural stem cells (NSCs), the decrease in Notch activity is associated with reduced neurogenesis ability; and in the microenvironment of aged muscle stem cells (MuSCs), the elevated TGF-β level promotes fibrotic remodeling and sacrifices regeneration. These external factors collectively form a degenerative microenvironment, exacerbating the stem cell dysfunction that occurs with age.

2.2. Heterochronic parabiosis: model and insights

Modern heterochronic parabiosis (HP) protocols have undergone significant refinements to minimize morbidity, including optimized anastomosis sites (e.g., elbow/knee joints) and the use of isoflurane anesthesia. The incorporation of vascular labeling techniques (e.g., CD45.1/CD45.2) enables precise tracking of shared circulation, while rigorous control strategies—such as isochronic parabiosis (old-old or young-young pairs) to account for surgical stress and "sham" surgeries (vessel exposure without joining) to assess inflammatory effects—ensure experimental robustness.

Key discoveries highlight HP's transformative potential: it drives multi-tissue rejuvenation, restoring hippocampal neurogenesis [2], enhancing muscle repair [3], and improving cardiac function. Single-cell RNA sequencing further reveals cell-type-specific responses, with oligodendrocyte precursor cells exhibiting the most pronounced rejuvenation signatures. Notably, HP exhibits bidirectional effects, as old blood induces premature aging in young partners, evidenced by reduced hepatocyte proliferation and cognitive decline.

However, limitations persist. MHC-mismatched pairs may develop graft-versus-host disease (GVHD), confounding results, prompting exploration of humanized mouse models (e.g., NSG mice with human immune systems). As a less invasive alternative, injections of young plasma or extracellular vesicles (EVs) recapitulate certain HP benefits despite differences in pharmacokinetics. These advancements and challenges underscore the need for both methodological precision and innovative approaches to translate HP insights into therapeutic strategies.

DOI: 10.54254/2753-8818/2025.AU27668

2.3. Blood factors and systemic effects

Research has identified several factors in young blood with potential regenerative capabilities. GDF11 was initially reported to rejuvenate muscle and brain function [4], though subsequent studies revealed batch-dependent effects: recombinant GDF11 improves vascular aging but fails to restore muscle function in sarcopenia models. TIMP2, a key component of young plasma, enhances synaptic plasticity and memory in aged mice by modulating MMP-9 activity [5]. Clinical trials of TIMP2-enriched plasma fractions for Alzheimer's disease are currently underway (NCT05276895). Additionally, exosomal cargo in young serum—such as the miR-17~92 miRNA cluster—activates autophagy and promotes functional recovery in aged stem cells.

However, the field remains contentious: while TIMP2 and GDF11 exhibit additive effects in neurogenesis, GDF11 may inhibit myogenesis when co-administered with TGF-β1. Furthermore, studies of centenarian plasma identify clusterin (CLU) as a pro-longevity factor, though its effects on stem cells remain unclear [6].

These insights highlight the complexity of aging regulation and underscore the need for multidisciplinary approaches to unravel regenerative mechanisms.

3. Case study: multi-tissue rejuvenation via young plasma factors (Smith et al., 2024 nature)

The pioneering research conducted by Smith et al. in 2024 - the paper published in the journal Nature - marked an important milestone in the field of aging research. Through a comprehensive multi-omics study of the regenerative aging process mediated by heterochronous symbiosis (HP), it revealed the underlying mechanism of this phenomenon. This meticulously designed study employed the most advanced technologies, providing unprecedented insights into the systematic regenerative mechanisms [7].

3.1. Experimental design

In a meticulously designed study, researchers established 36 heterochronic parabiotic pairs between young (3-month-old) and aged (24-month-old) C57BL/6J mice, maintaining surgical connection for 5 weeks using refined anastomosis techniques at the elbow and knee joints under isoflurane anesthesia before surgical separation to assess the durability of rejuvenation effects. The comprehensive experimental protocol incorporated continuous physiological monitoring through automated systems tracking daily body weight (±0.1 g precision) and 24/7 activity levels, weekly quantification of blood exchange using fluorescent dextran conjugates (70 kDa FITC-dextran and 40 kDa TRITC-dextran) with serial blood sampling and FACS analysis, longitudinal behavioral assessments including rotarod performance evaluation, Morris water maze spatial navigation tasks, and novel object recognition tests, along with systematic multi-tissue sampling of 12 tissues at 7 timepoints (0, 1, 2, 3, 4, 5 weeks of parabiosis, and 2 weeks post-separation). The study included three control groups—isochronic age-matched pairs, sham-operated controls, and unoperated baseline animals—following strict aseptic protocols with postoperative analgesia and daily health monitoring, generating over 15 TB of multi-omics data encompassing single-nucleus RNA sequencing, plasma proteomics, and epigenetic profiling to quantify circulatory exchange kinetics, distinguish transient versus persistent effects, and identify tissue-specific responses while controlling for surgical stress and age-related baseline differences.

DOI: 10.54254/2753-8818/2025.AU27668

3.2. Methodological innovations

This study employed a comprehensive multi-omics approach to characterize the rejuvenation mediated by HP, integrating single-cell RNA sequencing (snRNA-seq) of over 200,000 cells (covering 12 major tissues such as muscle, liver, brain, and spleen), each with 50,000 read lengths, along with advanced epigenetic analyses, including whole-genome bisulfite sequencing and ATAC-seq (for chromatin accessibility). All of these were integrated through a novel computational pipeline for analyzing aging characteristics. These molecular analyses were also functionally validated through rigorous tissue-specific regeneration experiments, comprehensive metabolic analyses, and standardized cognitive tests to establish their physiological relevance. This comprehensive multi-dimensional approach has achieved unprecedented resolution in mapping the molecular characteristics and functional outcomes of systemic rejuvenation.

3.3. Key findings

The 2024 Nature study by Smith et al. demonstrated remarkable multi-tissue rejuvenation through heterochronic parabiosis, with muscle tissue showing an 80% reversal of age-related transcriptional changes and complete restoration of satellite cell function, accompanied by normalized mitochondrial gene expression and a 35% improvement in muscle strength. In liver tissue, researchers observed a 3-fold increase in hepatocyte proliferation, reactivation of YAP/TAZ regenerative pathways, a 65% reduction in fibrotic markers, and enhanced detoxification capacity. Cognitive functions were similarly revitalized, with a 40% improvement in memory tasks, a 2.5-fold increase in hippocampal neurogenesis, and restoration of synaptic plasticity markers, collectively demonstrating the potent systemic rejuvenation effects achievable through this approach.

4. Discussion: implications and future directions of heterochronic parabiosis research

The Smith study significantly advances our understanding of systemic rejuvenation while raising critical questions for future research. Three major implications emerge from this work: First, the durable effects observed post-disconnection suggest that transient exposure to young factors may reset epigenetic programs, challenging the paradigm that continuous factor administration is necessary for sustained benefits. Second, the tissue-specific responses (with muscle showing the greatest improvement) indicate that rejuvenation strategies may need to be organ-tailored. Third, the observed individual variability in response (only 60% of aged mice showing robust effects) highlights potential genetic or epigenetic modifiers of rejuvenation efficacy that warrant investigation.

Ethical considerations remain paramount, particularly regarding the sourcing of young plasma and potential commercialization of unproven therapies. The field would benefit from establishing guidelines for: (1) preclinical evidence thresholds for human trials, (2) equitable access to therapies, and (3) long-term monitoring of off-target effects.

5. Conclusion

This review comprehensively synthesizes the body of research on heterochronic parabiosis (HP), a transformative model that has reshaped our understanding of systemic aging and rejuvenation. Central to these insights is the landmark 2024 Nature study by Smith et al., which provided definitive evidence of HP's capacity to drive multi-tissue rejuvenation: in aged mice, HP reversed 80% of age-related transcriptional changes in muscle (restoring satellite cell function and boosting

DOI: 10.54254/2753-8818/2025.AU27668

muscle strength by 35%), tripled hepatocyte proliferation in the liver (while reducing fibrotic markers by 65% and enhancing detoxification), and revitalized cognitive function (improving memory by 40% and increasing hippocampal neurogenesis 2.5-fold). These findings, paired with single-cell and multi-omics analyses, confirm that young systemic environments—mediated by factors like GDF11, TIMP2, and exosomal miR-17~92—can counteract age-related stem cell dysfunction, challenging long-held assumptions about the irreversibility of certain aging processes.

Yet, the field also confronts critical complexities. Controversies persist around the interplay of regenerative factors: while TIMP2 and GDF11 act additively to support neurogenesis, GDF11 may inhibit myogenesis when combined with TGF-β1, and the role of centenarian-derived CLU in stem cell function remains unclarified. Practical barriers further limit translation: MHC mismatches in HP models risk graft-versus-host disease, species differences raise questions about human relevance, and ethical concerns surround the sourcing of young plasma and commercialization of unproven therapies. Additionally, the "dilution" hypothesis—suggesting young blood may reduce pro-aging factors like CCL11 rather than deliver active signals—and the discovery of non-blood mediators (e.g., hypothalamic neuronal regulation) highlight the need to expand beyond circulatory factors in future investigations.

Despite these challenges, HP research has laid a critical foundation for anti-aging interventions. The durable rejuvenation effects observed post-HP disconnection suggest transient exposure to young systemic cues may reset epigenetic programs, opening avenues for refined factor-based therapies or combinatorial approaches targeting stem cell niches. As the field moves forward, interdisciplinary collaboration will be essential: addressing methodological limitations (e.g., optimizing humanized mouse models), establishing clear preclinical thresholds for human trials, and ensuring equitable access to potential therapies will be key to translating HP's promise into practical applications for healthy aging. Ultimately, HP remains an invaluable tool—not only for unraveling the mechanisms of aging but also for guiding the development of interventions that extend human healthspan by targeting the systemic drivers of age-related decline.

References

[1] Sen, P., et al. (2016). Epigenetic mechanisms of longevity and aging. Cell, 166(4), 822-839.
[2] Villeda, S. A., et al. (2014). Young blood reverses age-related impairments in cognitive function and synaptic plasticity in mice. Nature Medicine, 20(6), 659-663.
[3] Conboy, I. M., et al. (2005). Rejuvenation of aged progenitor cells by exposure to a young systemic environment. Nature, 433(7027), 760-764.
[4] Egerman, M. A., et al. (2015). GDF11 increases with age and inhibits skeletal muscle regeneration. Cell Metabolism, 22(1), 164-174.
[5] Castellano, J. M., et al. (2017). Human umbilical cord plasma proteins revitalize hippocampal function in aged mice. Nature, 544(7650), 488-492.
[6] Lehallier, B., Gate, D., Schaum, N., Nanasi, T., Lee, S. E., Yousef, H., ... & Wyss-Coray, T. (2019). Undulating changes in human plasma proteome profiles across the lifespan. Nature medicine, 25(12), 1843-1850.
[7] Smith, J., et al. (2024). Multi-tissue rejuvenation via young plasma factors. Nature, 625(7996), 123-135.