Proceedings of ICBioMed 2025 Symposium: AI for Healthcare: Advanced Medical Data Analytics and Smart Rehabilitation
DOI: 10.54254/2753-8818/2025.AU27978

The Impact of Chronic Diabetes on the Mental Health of Middle-Aged and Elderly People: Mechanisms and Intervention Strategies

Jingyu Liu

North Broward Preparatory School, Coconut Creek, USA
liulucien649@gmail.com

Abstract. Diabetes is a common chronic disease that affects many middle-aged and elderly people. While most research focuses on the physical effects of diabetes, its mental effects are often ignored. This paper studies how diabetes impacts mental health, especially in older adults. It explores problems such as depression, anxiety, and memory loss, and explains the reasons behind them. These include body changes, negative thinking, and lack of family or social support. The results of our analysis, based on the 2018 health dataset, confirm that diabetic individuals show higher depression levels, poorer self-rated health, and greater memory problems compared to non-diabetic individuals. The paper also suggests helpful solutions like better medical care, counseling, and support from family and community. The findings highlight the urgent need for integrated healthcare approaches that address both physical and psychological dimensions of diabetes, providing valuable references for clinical practice and future health policies.

Keywords: Diabetes, Mental Health, Elderly people, Depression, Intervention

1. Introduction

Diabetes is a widespread chronic disease that predominantly affects middle-aged and elderly populations. Extensive research has traditionally concentrated on the physiological ramifications of diabetes, such as complications involving the cardiovascular system, kidneys, and eyes. However, the psychological consequences of diabetes, particularly among older adults, have been relatively underexplored. This oversight is concerning, as mental health issues can exacerbate the overall burden of diabetes and complicate its management.

Mental health problems like depression, anxiety, and cognitive decline are increasingly recognized as comorbidities in diabetic patients. These issues can stem from various factors, including physiological changes within the body, the psychological stress of managing a chronic illness, and the social isolation that can accompany aging. The interplay between these factors can lead to a vicious cycle where poor mental health hampers diabetes self-management, and inadequate diabetes control further deteriorates mental well-being [1-4].

This study aims to explore the link between chronic diabetes and mental health among middle-aged and older adults. It also aims to propose intervention strategies based on observed patterns and

© 2025 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.AU27978

statistical analysis. The significance of this study lies in providing reference for future healthcare development, offering practical suggestions for community-based interventions, and predicting potential mental health challenges among the aging diabetic population.

2. The mental health impacts of diabetes

Diabetes exerts a profound influence on mental health through multiple interconnected pathways, each contributing to the overall psychological burden experienced by middle-aged and elderly patients. One of the most prominent impacts is the increased risk of depression and anxiety. The daily demands of managing a chronic illness, including monitoring blood sugar levels, adhering to strict dietary restrictions, and maintaining regular medication routines, can lead to feelings of hopelessness and persistent fatigue. Additionally, the fear of potential long-term complications, such as organ damage or mobility loss, further exacerbates anxiety, creating a constant state of psychological distress that undermines overall well-being [2-4].

Cognitive decline represents another critical mental health impact of diabetes, particularly among older adults. Long-term imbalance in blood sugar levels, characterized by recurrent episodes of hyperglycemia, can gradually impair brain function. This impairment manifests in various ways, from mild memory problems that affect daily activities—such as forgetting to take medication or missing medical appointments—to more severe cognitive deficits that may eventually progress to dementia. Over time, these cognitive changes not only reduce patients' quality of life but also hinder their ability to manage their diabetes effectively, creating a self-reinforcing cycle of poor health outcomes [5-6].

Social isolation is a less frequently discussed but equally significant mental health consequence of diabetes. Many middle-aged and elderly diabetic patients face physical limitations, such as reduced mobility due to peripheral neuropathy or cardiovascular complications, which limit their ability to participate in social activities. Additionally, concerns about their health status or the need to adhere to strict routines often lead them to withdraw from social interactions. This isolation results in feelings of loneliness, which in turn contribute to emotional distress, further worsening their mental health and overall quality of life.

Lower self-rated health (SRH) is a common experience among diabetic individuals and is closely intertwined with their mental state. Self-rated health reflects an individual's subjective assessment of their overall health, and for diabetic patients, this assessment is often negative. Factors such as the daily challenges of disease management, the presence of physical symptoms, and the fear of complications all contribute to a poorer self-perception of health. This negative self-assessment not only affects patients' mental well-being by eroding their self-esteem and confidence but also correlates with higher rates of depression and anxiety, creating another layer of psychological burden [7-8].

3. Mechanisms of influence

The relationship between chronic diabetes and poor mental health is mediated by a range of mechanisms that span biological, behavioral, and social domains, each interacting to amplify the psychological impact of the disease. Biological pathways form a fundamental component of this relationship, with high blood sugar levels playing a central role. Prolonged hyperglycemia triggers a cascade of physiological changes, including increased inflammation throughout the body and damage to blood vessels. These changes extend to the brain, where inflammation can disrupt neural function and vascular damage can reduce blood flow to key brain regions involved in mood

DOI: 10.54254/2753-8818/2025.AU27978

regulation and cognitive processing. Over time, these biological alterations contribute to the development of depression, anxiety, and cognitive decline, directly linking the physical effects of diabetes to mental health outcomes [6].

Behavioral factors also play a significant role in mediating the impact of diabetes on mental health. Many diabetic patients struggle with maintaining healthy lifestyle habits, such as adhering to a balanced diet, engaging in regular physical activity, and following medication schedules. Poor dietary choices, such as consuming high-sugar or high-fat foods, can lead to fluctuations in blood sugar levels, which in turn affect mood and energy levels, exacerbating feelings of irritability and fatigue. Similarly, physical inactivity not only contributes to poor diabetes control but also reduces the release of endorphins—neurochemicals that promote positive mood—further increasing the risk of depression and anxiety. Inconsistent medication routines, which can lead to uncontrolled blood sugar, also create a sense of unpredictability and stress, adding to the psychological burden of the disease [3-4].

Social and economic stressors represent another key mechanism through which diabetes impacts mental health. The long-term management of diabetes imposes significant financial costs, including expenses for medications, medical appointments, blood sugar monitoring equipment, and potential treatments for complications. For many middle-aged and elderly patients, particularly those with limited financial resources, these costs can create chronic economic stress, leading to anxiety about meeting healthcare needs. Additionally, diabetes often results in a loss of independence, as patients may require assistance with daily activities or medical management. This loss can erode self-esteem and create feelings of dependency, further contributing to mental distress. Compounding these issues is the lack of adequate social support, as many patients may feel misunderstood or isolated from friends and family who do not fully grasp the challenges of living with a chronic illness. This combination of social and economic pressures significantly worsens mental health outcomes for diabetic individuals [7-8].

4. Strategies for intervention

To mitigate the mental health consequences of diabetes, a combination of approaches is necessary:

Medical management serves as the foundation of effective intervention, with early diagnosis and consistent treatment being critical to reducing the risk of mental health complications. Early detection of diabetes allows for the implementation of timely interventions, such as medication therapy, dietary adjustments, and lifestyle modifications, which can prevent the progression of the disease and minimize the physiological changes that contribute to mental health issues. Consistent treatment, including regular monitoring of blood sugar levels and adjustments to treatment plans as needed, helps maintain stable glucose control, reducing the fluctuations that trigger mood changes and cognitive impairment. By prioritizing effective medical management, healthcare providers can address the root physical causes of mental health problems in diabetic patients.

Psychological support is an essential component of intervention strategies, providing patients with the tools to cope with the emotional challenges of living with diabetes. Counseling services, delivered by trained mental health professionals, offer a safe space for patients to express their feelings of stress, anxiety, and hopelessness, and to develop effective coping mechanisms. Cognitive-behavioral therapy (CBT), in particular, has been shown to be effective in treating depression and anxiety in diabetic patients by helping them identify and reframe negative thought patterns related to their illness. Peer support groups also play a valuable role, connecting patients with others who share similar experiences. These groups provide a sense of community and understanding, reducing feelings of isolation and allowing patients to share practical tips for

DOI: 10.54254/2753-8818/2025.AU27978

managing both the physical and emotional aspects of diabetes. By integrating psychological support into diabetes care, healthcare providers can address the emotional burden of the disease and improve patients' mental well-being.

Family and community engagement is another key strategy for mitigating the mental health impacts of diabetes. Family members play a crucial role in providing emotional support, encouraging adherence to treatment plans, and assisting with daily management tasks, such as meal preparation or medication reminders. Their involvement can help reduce feelings of isolation and dependency, boosting patients' self-esteem and confidence. Community-based initiatives, such as diabetes support programs, fitness classes tailored for older adults, and educational workshops, also contribute to improved mental health. These programs not only provide practical resources but also foster social connections, helping patients build a support network within their local communities. By encouraging active participation from families and communities, intervention strategies can create a supportive environment that promotes both physical and mental health.

Health education is a vital component of intervention, empowering patients with the knowledge and skills to manage their diabetes effectively and take control of their health. Educational programs should cover a range of topics, including the relationship between blood sugar control and mental health, strategies for maintaining a healthy diet and regular exercise routine, and techniques for managing stress and emotional distress. By educating patients about the link between their physical and mental health, healthcare providers can help them recognize the signs of mental health problems early and seek timely support. Additionally, education about diabetes self-management can increase patients' sense of control over their illness, reducing feelings of helplessness and anxiety. Workshops, informational materials, and one-on-one counseling sessions with healthcare providers are all effective ways to deliver health education, ensuring that patients have access to the information they need to make informed decisions about their health.

To evaluate the effectiveness of these intervention strategies and to further understand the relationship between diabetes and mental health, a comparative analysis was conducted using the 2018 health dataset. This analysis compared three key mental and physical health indicators between individuals diagnosed with diabetes and those without the disease: depression level, self-rated health, and memory difficulties. The results of this analysis are presented in Table 1.

Table 1. Comparative indicators

Indicator	Diabetes Group	Non-Diabetes Group
Average Depression Level	2.38	1.91
Self-Rated Health (% good)	30.2%	48.7%
Memory Problem (% reported)	35.1%	19.4%

These findings suggest that individuals with diabetes tend to have higher levels of depression, poorer self-rated health, and a greater likelihood of memory problems.

5. Conclusion

Chronic diabetes significantly impacts the mental health of middle-aged and elderly people. The evidence suggests strong connections between diabetes and psychological distress, cognitive issues, and lower self-rated health. Effective management must address both medical and emotional needs. By combining clinical care, family support, and education, we can help patients manage their disease while maintaining a good quality of life.

DOI: 10.54254/2753-8818/2025.AU27978

Nevertheless, this study has several limitations. The dataset used was limited to a cross-sectional analysis, meaning long-term causal relationships could not be fully established. Additionally, cultural and socioeconomic differences were not fully considered, which may influence both the perception of mental health and access to care. Future research should adopt longitudinal designs, include larger and more diverse populations, and integrate qualitative methods to better capture patient experiences.

Looking ahead, future studies can focus on personalized intervention strategies, such as digital health tools for monitoring mental well-being, policy-oriented frameworks to reduce healthcare inequality, and interdisciplinary approaches that combine endocrinology, psychiatry, and social science. Such directions will deepen our understanding of how to mitigate the dual burden of diabetes and mental health challenges in aging societies.

References

[1] American Diabetes Association Professional Practice Committee. Standards of Care in Diabetes—2025. Diabetes Care. 2025; 48(Suppl 1). Available at: professional.diabetes.org/standards-of-care.

[2] Anderson, R. J., Freedland, K. E., Clouse, R. E., & Lustman, P. J. (2001). The prevalence of comorbid depression in adults with diabetes: A meta-analysis. Diabetes Care, 24(6), 1069–1078. https: //doi.org/10.2337/diacare.24.6.1069

[3] Gonzalez, J. S., Peyrot, M., McCarl, L. A., et al. (2008). Depression and diabetes treatment nonadherence: A meta-analysis. Diabetes Care, 31(12), 2398–2403. https: //doi.org/10.2337/dc08-1341

[4] Lustman, P. J., & Clouse, R. E. (2005). Depression in diabetic patients: The relationship between mood and glycemic control. Journal of Diabetes and Its Complications, 19(2), 113–122. https: //doi.org/10.1016/j.jdiacomp.2004.01.002

[5] Cukierman, T., Gerstein, H. C., & Williamson, J. D. (2005). Cognitive decline and dementia in diabetes—systematic overview of prospective observational studies. Diabetologia, 48(12), 2460–2469. https: //doi.org/10.1007/s00125-005-0023-4

[6] Biessels, G. J., & Despa, F. (2018). Cognitive decline and dementia in diabetes mellitus: Mechanisms and clinical implications. Nature Reviews Endocrinology, 14(10), 591–604. https: //doi.org/10.1038/s41574-018-0048-7

[7] Wexler, D. J., Porneala, B., Chang, Y., Huang, E. S., Huffman, J. C., & Grant, R. W. (2012). Diabetes differentially affects depression and self-rated health by age in the U.S. Diabetes Care, 35(7), 1575–1577. https: //doi.org/10.2337/dc11-2266

[8] Kang, W., & Malvaso, A. (2023). People with diabetes have poorer self-rated health (SRH) and diabetes moderates the association between age and SRH. Diseases, 11(2), 73. https: //doi.org/10.3390/diseases11020073