Proceedings of the 6th International Conference on Biological Engineering and Medical Science
DOI: 10.54254/2753-8818/2026.32734

The Chemistry of Food Preservatives

Chenxi Dai

School of Applied Chemistry, Xi'an Jiaotong-Liverpool University, Suzhou, China
Chenxi.Dai2402@student.xjtlu.edu.cn

Abstract. Food preservatives play an indispensable role in contemporary food processing and distribution systems by inhibiting microbial proliferation and retarding physicochemical deterioration, thereby ensuring food safety and extending product shelf life. They are extensively utilized in diverse food matrices, including meat products, dairy systems, and beverages. With increasing regulatory scrutiny and consumer demand for higher safety standards, the scientific optimization of preservative formulations has become critical for enhancing antimicrobial efficacy while minimizing additive levels and maintaining product stability. This study systematically examines the mechanisms of commonly used food preservatives, with particular emphasis on the functional principles of organic acids and antioxidants within complex food systems. The findings indicate that preservative efficacy is strongly influenced by intrinsic food properties and environmental parameters, particularly pH and formulation ratios.

Keywords: Food, Microorganism, Spoilage, Acid, Preservative

1. Introduction

Food stability remains a central concern in modern food science, as complex chemical and biochemical reactions continuously compromise product quality, safety, and shelf life. Deterioration processes arise not only from microbial proliferation but also from intrinsic chemical instabilities within food matrices, including oxidation, non-enzymatic browning, and hydrolytic degradation. A systematic understanding of the chemical principles governing both spoilage and preservation is therefore essential for advancing rational formulation strategies. This study addresses three interrelated dimensions of the topic. First, it elucidates the chemical basis of food spoilage by examining microbial metabolism alongside non-microbial reaction pathways that drive physicochemical deterioration. Second, it analyzes the molecular mechanisms underlying mainstream food preservatives, particularly organic acids and antioxidant systems, with emphasis on their physicochemical interactions within food environments. Third, it evaluates the key chemical factors influencing preservative efficacy, including pH, water activity, and compositional dynamics. A structured literature review and analytical synthesis of authoritative academic sources were employed to construct a coherent theoretical framework. By integrating spoilage chemistry with preservation mechanisms, this study establishes a physicochemical perspective that supports evidence-based preservative design and promotes.

© 2026 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2026.32734

2. Chemical basis of food spoilage

2.1. Chemical mechanisms of microbial spoilage

Microbial spoilage of food results from the metabolic activities of bacteria, yeasts, and molds that utilize food components as substrates for growth and reproduction. During proliferation, spoilage microorganisms secrete a suite of extracellular enzymes such as proteases, lipases, and carbohydrate-degrading hydrolases that catalyze the breakdown of macromolecules into low-molecular-weight products. These enzyme-mediated transformations represent essential biochemical pathways by which cellular metabolism accelerates quality loss.

Physicochemical conditions within the food matrix strongly modulate microbial metabolism. The pH influences both the enzyme catalytic efficiency and the growth range of microorganisms: most spoilage bacteria prefer a near-neutral pH range, whereas many fungi can tolerate broader acidic conditions, resulting in differential spoilage patterns across food types [1]. Water activity (aw) is another critical determinant, as sufficient free water is required for enzymatic function, nutrient diffusion, and cellular transport; reduction of aw through drying or solute addition limits microbial proliferation by restricting biochemical mobility [1]. Intracellular metabolic reactions also produce organic acids and gases that further alter local pH and exacerbate chemical deterioration. Therefore, microbial spoilage embodies a complex interplay between biochemical metabolism and the surrounding chemical environment.

2.2. Chemical mechanisms of non-microbial spoilage

A predominant pathway is lipid oxidation, a free radical chain reaction initiated by hydrogen abstraction from unsaturated fatty acids, followed by molecular oxygen addition and propagation to lipid hydroperoxides; the subsequent decomposition of these hydroperoxides yields aldehydes, ketones, and short-chain acids responsible for off-flavors and loss of nutritional quality [2]. Another critical non-enzymatic process is the Maillard reaction, which involves the condensation between reducing sugars and amino groups of proteins or amino acids, forming a complex series of intermediates and heterocyclic compounds that result in browning, flavor changes, and formation of advanced glycation end products [3]. Light-induced oxidation also accelerates deterioration, as photo-oxidative pathways generate excited singlet oxygen and reactive radicals that attack unsaturated lipids and pigments. Hydrolytic reactions, particularly triglyceride hydrolysis, contribute to the accumulation of free fatty acids under high moisture and elevated temperature conditions, further enhancing susceptibility to oxidation. Collectively, these non-microbial pathways reflect fundamental chemical instability in food matrices, governed by intrinsic composition and environmental factors.

3. Mainstream food preservatives

3.1. Antimicrobial mechanisms based on acid–base and membrane interactions

Organic acid preservatives constitute one of the most extensively applied categories of chemical preservatives in the food industry, including benzoic acid, sorbic acid, propionic acid, and lactic acid, all of which are weak carboxylic acids. When the environmental pH is lower than their respective pKa values, the proportion of the undissociated form (HA) increases. In this state, the molecule exhibits greater lipophilicity and is capable of penetrating microbial cell membranes via passive diffusion. Once inside the near-neutral cytoplasmic environment, the acid dissociates,

DOI: 10.54254/2753-8818/2026.32734

releasing protons (H^+) that reduce intracellular pH and disrupt the proton motive force and energy metabolism, thereby inhibiting cellular growth and proliferation. Furthermore, the corresponding anionic forms may interfere with key enzymatic systems and the tricarboxylic acid (TCA) cycle, aggravating metabolic imbalance [4].

For instance, benzoic acid, with a relatively lower pKa, is more suitable for acidic food systems; sorbic acid, containing conjugated double bonds, demonstrates strong antifungal activity; whereas propionic acid exhibits pronounced anti-mold effects in bakery products. Therefore, the rational selection of specific organic acids must consider food pH conditions and target microorganisms.

3.2. Nitrite preservatives and emerging preservative systems

Nitrite preservatives, particularly sodium nitrite, exert antimicrobial activity through the generation of reactive nitrogen species under acidic conditions. In meat systems, nitrite (NO_2^-) is protonated to form nitrous acid (HNO_2), which subsequently decomposes to yield nitric oxide (NO), nitrogen dioxide (NO_2), and dinitrogen trioxide (N_2O_3). These intermediates participate in nitrosation and redox reactions with microbial cellular constituents. Nitric oxide readily binds to iron–sulfur clusters and heme-containing enzymes, thereby inhibiting cytochrome-mediated electron transport and disrupting ATP synthesis. Concurrently, nitrosative stress interferes with spore germination and vegetative growth of Clostridium botulinum, which underlies the preservative efficacy of nitrite in cured meat products [5].

Nevertheless, excessive nitrite may react with secondary amines via nitrosation pathways, particularly under acidic or thermal conditions, leading to the formation of N-nitrosamines with recognized carcinogenic potential [6]. This duality necessitates strict regulatory control to balance antimicrobial effectiveness and toxicological safety. To reduce reliance on nitrites, emerging preservative systems—including bacteriocins (e.g., nisin), plant-derived polyphenols, and antimicrobial edible coatings—have been developed.

4. Factors affecting preservative efficacy

4.1. Environmental factors

The preservative efficacy of food systems is fundamentally determined by environmental physicochemical parameters, including temperature, water activity (aw), pH, and osmotic pressure.In contrast, low-temperature storage decreases molecular mobility and kinetic energy, effectively reducing reaction rates and microbial proliferation. Water activity represents the thermodynamic availability of free water required for microbial growth; most spoilage and pathogenic bacteria require aw values above 0.90. Consequently, reducing aw through dehydration or solute incorporation shifts the system beyond the microbial growth threshold and enhances preservation stability.

pH plays a critical role in both microbial physiology and the acid–base dissociation equilibria of weak-acid preservatives. Lower pH values increase the proportion of undissociated acid species, thereby promoting membrane permeation and intracellular acidification. Osmotic pressure in hypertonic environments, such as high-salt or high-sugar systems, induces plasmolysis and disrupts cellular osmotic homeostasis.

DOI: 10.54254/2753-8818/2026.32734

4.2. Food type factors

Variations in spoilage rates among different food categories are fundamentally determined by intrinsic matrix composition and structural properties. In contrast, commodities with reduced water activity, limited nutrient bioavailability, or compact structural matrices—such as dried cereals or high-sugar products—exhibit inherently lower microbial growth potential and enhanced storage stability. The physicochemical characteristics of the matrix therefore modulate reaction rate constants, diffusion behavior, and metabolic accessibility.

Preservative efficacy is further influenced by molecular interactions between antimicrobial agents and food components. Weak-acid preservatives may undergo phase partitioning between lipid and aqueous domains, reversible binding to proteins, or partial neutralization due to matrix buffering capacity, thereby altering their effective concentration in the aqueous phase where microbial activity predominates. Distribution coefficients and binding equilibria consequently determine antimicrobial availability. Additionally, processing technologies—including thermal sterilization, dehydration, fermentation, salting, and modified atmosphere packaging—extend shelf life by reducing initial microbial load, lowering water activity, modifying pH, and limiting oxygen-dependent reactions. Collectively, intrinsic matrix properties and processing-induced modifications govern spoilage kinetics and preservation efficiency in diverse food systems.

5. Conclusion

Food deterioration arises from the interplay between microbial metabolic activity and non-microbial chemical transformations, including lipid oxidation, hydrolysis, and enzymatic catalysis, all of which are fundamentally regulated by thermodynamic feasibility and kinetic parameters. Widely used preservatives such as organic acids and nitrites exert antimicrobial activity through acid–base dissociation equilibria, membrane permeability modulation, and redox- or nitrosation-mediated biochemical interference, thereby disrupting cellular homeostasis and metabolic flux. Their preservative performance is strongly modulated by environmental variables—including temperature-dependent reaction kinetics, water activity–controlled thermodynamic availability, pH-regulated dissociation equilibria, and osmotic stress—as well as intrinsic food matrix characteristics that influence molecular partitioning behavior, binding equilibria, and effective antimicrobial concentration within multiphase systems. These findings indicate that food preservation should be interpreted as a problem of physicochemical regulation within complex biological matrices rather than the mere addition of antimicrobial agents. The development of natural bioactive compounds, antimicrobial peptides, controlled-release delivery systems, and intelligent packaging technologies, combined with kinetic modeling and systems-level predictive analysis, may enable more precise and sustainable control of spoilage processes. Mechanism-guided, safety-oriented, and environmentally compatible preservative systems will therefore remain a central direction in contemporary food chemistry research.

References

[1] Chen, L. (2025). Changes in microbial metabolites during food mold growth and their effects on food quality. Industrial Microbiology, 55(3), 198–200.
[2] Ge, Y. T., Kou, B. Y., & Zhang, C. Y. (2025). Research progress on the application of natural antioxidants in meat product processing. Journal of Xinyang Normal University (Natural Science Edition), 38(4), 497-505.
[3] Lv, S. Y., Jiang, C. C., Yang, Y. C., Qiu, W. Q., Bao, J. Q., & Jin, Y. Z. (2026). Effects of Maillard reaction on physicochemical properties and flavor characteristics of hydrolysate from melon whelk meat. Food and

DOI: 10.54254/2753-8818/2026.32734

Fermentation Industries, 1-12. https: //doi.org/10.13995/j.cnki.11-1802/ts.045421.

[4] Qianjin Liu, Tiantian Li, Xiaonan Gong, Li Yang, Yaoguang Li, Biao Chen, Shengtao Ma, Jingnan Zhang, Shuai Wan, Jie Zhang, Lili Cai, Zhuo Wen, Junsong Zhang & Xiangyu Zang. (2025). Microbial community succession during storage of tobacco flavor liquid and the inhibitory effect of potassium sorbate on spoilage bacteria. Tobacco Science & Technology, 1-15.

[5] Suyan Zhang & Aiwu Gao. (2016). Research progress on the application of nitrite substitutes in meat products. Meat Research, 30(10), 45–48.

[6] Beibei Geng & Hongjun Zuo. (2025). Investigation and Analysis of a Food Poisoning Incident Caused by Misuse of Nitrite in Catering Establishments. Food Safety Guide, 19(30), 11-13.