Proceedings of CONF-APMM 2025 Symposium: Multi-Qubit Quantum Communication for Image Transmission over Error Prone Channels
DOI: 10.54254/2753-8818/2025.GL33588

Algebraic and Topological Approaches to Bloch's Theorem, Vortex Circulation, and Electrodynamics via Complex Analysis

Junyu Luo[1*], Zicheng Zhuang[2], Jiarui Guo[3]

[1] *Department of Physics, Hubei Normal University, Huangshi, China*
[2] *Department of Physics, Suzhou North America High School, Suzhou, China*
[3] *Department of Physics, Ulink College Guangzhou, Shenzhen, China*
[*] *Corresponding author: 18173957783@163.com*

Abstract. We present a unified algebraic–geometric framework in which three cornerstone problems in mathematical physics are interlaced through the lens of complex analysis and group theory. First, Bloch's theorem is recast as the classification of one-dimensional unitary characters of the Bravais translation group, whereby crystal momentum is naturally labeled by irreducible representations and spectral, transport, and topological properties of electronic bands are directly obtained, with contour-integral Green's-function constructions being employed to separate propagating and evanescent modes. Second, the classical residue theorem is reinterpreted within the topological language of π_1 and de Rham cohomology to demonstrate that conserved observables including planar electric flux and vortex circulation are shown to depend solely on local pole residues and loop winding numbers, thus bypassing coordinate integrals and exposing homotopy invariance in electrostatics and ideal fluid flows. Third, by means of a power-law conformal map, the electrostatic wedge problem is solved in closed form, the characteristic $1/r$ field singularity at sharp tips is thereby exposed, and the isomorphism $\mathrm{PSL}(2,\mathbb{C}) \simeq \mathrm{SO}^{+}(1,3)$ is subsequently unveiled, identifying Lorentz transformations with Möbius maps on the celestial sphere. Finally, this synthesis is shown to deepen the interplay between characters, residues, and analytic mappings and to lay the conceptual groundwork for extensions to nonlinear field theories, singular Riemann-surface methods, and modern celestial–CFT approaches to scattering amplitudes.

Keywords: Bloch's theorem, Residue theorem, Conformal mapping, Complex analysis, Topology

1. Introduction

Complex analysis and group-representation theory have long been celebrated for their unifying power in addressing diverse problems across theoretical and applied physics [1–4]. In solid-state physics, Bloch's theorem is founded on the unitary representations of the crystal translation group, by which electronic wavefunctions are decomposed into plane-wave factors multiplied by lattice-periodic functions [5–7]. In fluid dynamics and electromagnetism, local singularities of analytic functions are related to global conservation laws, such as circulation around vortices or net charge, by the residue theorem, and the topological nature of these invariants is made clear via contour deformations [8, 9]. By

© 2026 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.GL33588

preserving angles under holomorphic transformations, complicated boundary geometries are transformed into canonical domains by conformal mapping, whereby Laplace-equation boundary-value problems that would otherwise be intractable are converted into standard, solvable configurations [10, 11]. A natural bridge between abstract mathematical invariants and concrete physical phenomena is provided by complex analysis. However, in much of the existing literature complex analysis is typically treated in isolation. For example, in the physics-oriented texts by Jackson, Arfken and Weber and Morse and Feshbach [3, 12, 13], almost exclusive attention is paid to step-by-step evaluation of contour integrals and residue calculations in specific problems, without ascending to the underlying abstract framework that would connect these computations to deeper mathematical principles such as the role of complex structures in revealing invariants and symmetry groups of physical systems. Conversely, in the more mathematically rigorous treatments by Ahlfors, Conway and Rudin [14–16], complex analysis and other branches of mathematics are developed axiomatically, with definitions, theorems and proofs presented in parallel, yet little effort is made to demonstrate how these abstract tools can be deployed in a unified, practical fashion to illuminate physical phenomena. As a result, the unifying role of complex analysis in linking mathematical invariants to physical problems, as well as the abstract mathematical perspective underlying these problems, has often remained underappreciated. In this review, previous efforts are built upon by refining the approach to understanding and solving physical problems through a unified mathematical–physical perspective. The core framework of complex analysis, enriched by insights from group representation theory and topological homotopy, is adopted. By juxtaposing three archetypal problems drawn from distinct areas of physics, the corresponding invariants in complex analysis are highlighted and the influence of local algebraic and topological structures on global behavior is demonstrated.

In Section 1, a streamlined derivation of Bloch's theorem is presented, rooted in the theory of one-dimensional unitary characters of a three-dimensional Bravais lattice. In this algebraic treatment, the reason why crystal momentum labels the irreducible representations of the translation group is clarified, and the periodic structure of eigenfunctions in reciprocal space is illuminated [5, 17]. In Section 2, a topological perspective on the residue theorem is offered: after it is framed in terms of de Rham cohomology and fundamental-group generators, the theorem is applied to compute the circulation around a point vortex in a two-dimensional incompressible fluid, and it is demonstrated that the circulation is entirely determined by the vortex strength and the contour's winding number [8, 18]. In Section 3, the power-law conformal map that carries a conducting wedge onto the upper half-plane is introduced; by solving the corresponding Dirichlet problem there and pulling back, explicit formulas for the electrostatic potential, field, and surface charge density in the original wedge are obtained, highlighting the singular field enhancement near sharp edges [10, 11]. Furthermore, not only is the solution of partial differential equations simplified by conformal mappings, but the algebraic structures underlying certain problems in physics are also revealed within the framework of complex analysis, thereby aiding the understanding of symmetries from a complex analysis perspective. Subsequently, the isomorphism between the Möbius transformation group and the proper orthochronous Lorentz group is introduced, and it is shown how four-vectors represented by Hermitian matrices present a complex analysis perspective for understanding the Lorentz transformation in special relativity. By algebraic, topological, and analytical methods being woven together, the role of complex analysis as a common thread linking seemingly disparate physical phenomena is illustrated by this work. A novel perspective based on complex analysis, focusing on invariants and symmetries, is provided for physicists and learners alike, in order to better understand the mathematical structures underlying physical laws and to offer more effective solutions to physical problems. Additionally, the practical application of complex analysis and its associated abstract theories in physical models is highlighted, contribut-

DOI: 10.54254/2753-8818/2025.GL33588

ing to a deeper understanding of the mathematical foundations among readers. In the following, each section will be proceeded through in turn, with the goal of leveraging complex analysis more deeply to uncover the pivotal role played by these invariants in both solving and interpreting the underlying physical problems.

2. Bloch's Theorem from the views of group-representation theory and complex analysis

2.1. Unitary representation of the translation group and bloch states

The interplay between symmetry and wave dynamics lies at the heart of modern condensed-matter physics. Consider the single-particle Hamiltonian

$$\hat{H} = -\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{r}), \tag{1}$$

which remains invariant under any Bravais-lattice translation $\mathbf{R} = n_1\mathbf{a}_1 + n_2\mathbf{a}_2 + n_3\mathbf{a}_3$ [7] . These translations form the Abelian group

$$G = \left\{\hat{T}_{\mathbf{R}} : \hat{T}_{\mathbf{R}}\psi(\mathbf{r}) = \psi(\mathbf{r}+\mathbf{R})\right\}, \tag{2}$$

and since $[\hat{H}, \hat{T}_{\mathbf{R}}] = 0$, one may seek simultaneous eigenfunctions:

$$\hat{H}\,\psi = E\,\psi, \qquad \hat{T}_{\mathbf{R}}\,\psi = C(\mathbf{R})\,\psi. \tag{3}$$

By the theory of one-dimensional unitary representations of an Abelian group, the character $C : G \to U(1)$ satisfies

$$C(\mathbf{R}_1 + \mathbf{R}_2) = C(\mathbf{R}_1)\,C(\mathbf{R}_2), \qquad |C(\mathbf{a}_i)| = 1, \tag{4}$$

and thus admits the parametrization

$$C(\mathbf{a}_i) = e^{i\alpha_i}, \quad \alpha_i \in \mathbb{R}. \tag{5}$$

Repeated application then gives

$$C(\mathbf{R}) = \exp\Big(i\sum_i n_i\alpha_i\Big), \tag{6}$$

and upon identifying $\alpha_i = \mathbf{k}\cdot\mathbf{a}_i$ one arrives at the crystal momentum label:

$$C(\mathbf{R}) = e^{i\mathbf{k}\cdot\mathbf{R}}. \tag{7}$$

Hence the Bloch eigenfunctions assume the familiar form

$$\psi_{\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}\,u_{\mathbf{k}}(\mathbf{r}), \qquad u_{\mathbf{k}}(\mathbf{r}+\mathbf{R}) = u_{\mathbf{k}}(\mathbf{r}). \tag{8}$$

Physically, the plane-wave factor $e^{i\mathbf{k}\cdot\mathbf{r}}$ encodes the long-range propagation of an electron with crystal momentum $\hbar\mathbf{k}$, while the periodic function $u_{\mathbf{k}}(\mathbf{r})$ demonstrates the translational symmetry of the lattice. Group representation theory not only reveals the algebraic structure and symmetry of physical laws, but also provides a foundation for specific physical calculations.Based on the group representation theory, we obtain k. With this, we can carry out various subsequent physical calculations. Let's take the Green function, which is an important part of the condensed matter theory, as an example.

DOI: 10.54254/2753-8818/2025.GL33588

2.2. The complex analysis method in Green's function

To probe spectral and spatial correlations, one introduces the retarded Green's function at energy E.First,we consider the one-dimensional situation

$$G(E;\mathbf{r},\mathbf{r}') = \left\langle \mathbf{r} \,\middle|\, (E - \hat{H} + i0^+)^{-1} \middle| \mathbf{r}' \right\rangle. \tag{9}$$

Using the completeness of the Bloch states, one may write

$$G(E;x,x') = \frac{1}{2\pi} \int_{-\pi/a}^{\pi/a} \frac{e^{\,ik(x-x')}}{E - \varepsilon_n(k) + i0^+} \,\mathrm{d}k. \tag{10}$$

By analytically continuing k into the complex plane and closing the contour $\mathcal{C}$ around all simple poles satisfying $E = \varepsilon_n(k)$, one obtains

$$G(E;x,x') = \frac{1}{2\pi i} \oint_{\mathcal{C}} \frac{e^{\,ik|x-x'|}}{E - \varepsilon_n(k)} \,\mathrm{d}k, \tag{11}$$

where $\mathcal{C}$ is taken counterclockwise around each pole. By residue calculus, the pole contributions determine spatial decay and oscillation in one dimension. In three dimensions, one integrates over complex surfaces in $\mathbf{k}$-space and sums band-branch residues, recovering the full lattice Green's function in 3D crystals [19, 20].

Equivalently, via spectral decomposition

$$G(E;\mathbf{r},\mathbf{r}') = \sum_{\alpha} \frac{\psi_\alpha(\mathbf{r})\, \psi_\alpha^*(\mathbf{r}')}{E - E_\alpha + i0^+}, \tag{12}$$

and setting $\mathbf{r}' = \mathbf{r}$, taking the imaginary part yields

$$\Im\, G(E;\mathbf{r},\mathbf{r}) = -\pi \sum_{\alpha} \left|\psi_\alpha(\mathbf{r})\right|^2 \delta(E - E_\alpha), \tag{13}$$

so that the local density of states is

$$\rho(E) = -\frac{1}{\pi} \Im\, G(E;\mathbf{r},\mathbf{r}) = \sum_{\alpha} \left|\psi_\alpha(\mathbf{r})\right|^2 \delta(E - E_\alpha), \tag{14}$$

directly corresponding to spectroscopic observables. This complex-analysis approach is seen to offer several key advantages. Methodologically, by promoting k to a complex variable and closing the integration contour around the simple poles defined by $E = \varepsilon_n(k)$, an explicit decomposition of the Green's function into contributions from individual Bloch branches is obtained. Not only is numerical evaluation thereby streamlined,since only a discrete set of poles need be located and summed over rather than sampling the entire real-k axis, but propagating modes are also naturally separated from evanescent or surface states (complex-k poles). Physically, the oscillation period and decay length of a given electronic mode are directly encoded by each residue term. In particular, bulk plane-wave propagation is signaled by real k_p, while exponentially decaying evanescent waves that govern tunneling and surface-state penetration depths are indicated when $\Im\, k_p \neq 0$. Moreover, a seamless generalization to multiband and anisotropic lattices is afforded by simply tracking additional poles from each branch and deforming the contour to respect branch cuts introduced by non-trivial band topology. Thus, not only is a physically transparent picture of spectral and spatial correlations in crystalline

DOI: 10.54254/2753-8818/2025.GL33588

media provided by the contour-integral Green's function, but a robust computational backbone for evaluating local density of states, scattering amplitudes, and interface phenomena in both conventional and topological materials is also formed. The k obtained through group representation theory is also served as the cornerstone of the band theory, and based on this, a series of calculations for transport properties can be conducted. Next, the calculation related to electron velocity and the effective mass tensor will be demonstrated as examples.

2.3. Transport quantities and effective mass

From the band dispersion $E_n(\mathbf{k})$ one defines the semiclassical group velocity and effective-mass tensor

$$\mathbf{v}_n(\mathbf{k}) = \frac{1}{\hbar}\,\nabla_{\mathbf{k}} E_n(\mathbf{k}), \tag{15}$$

$$\left[m^{-1}(\mathbf{k})\right]_{\alpha\beta} = \frac{1}{\hbar^2}\,\frac{\partial^2 E_n}{\partial k_\alpha\,\partial k_\beta}\,. \tag{16}$$

Here $\mathbf{v}_n(\mathbf{k})$The instantaneous velocity of a wave-packet centered at $\mathbf{k}$ is described, encoding how the electron responds to an applied field. The curvature of the band, captured by the second derivatives of $E_n(\mathbf{k})$, is naturally interpreted as an effective mass $m_{\alpha\beta}(\mathbf{k})$ that generally differs from the free-electron mass m. Physically, the inertia of carriers moving through the periodic potential is quantified by the effective mass. In regions of low curvature, since $\partial^2 E/\partial k^2$ is small, small eigenvalues are possessed by the effective-mass tensor, indicating sluggish carrier dynamics and low mobility. Conversely, in regions of high curvature, carriers are made to behave as if they were light and external forces are responded to quickly.

In many experimental contexts, such as cyclotron resonance or quantum oscillation analysis, a scalar effective mass m^* is commonly used to simplify the description. This scalar m^* can be viewed as the averaged value or the trace of the tensor $m_{\alpha\beta}$. For isotropic systems, a common definition is

$$\frac{1}{m^*} = \frac{1}{d}\,\mathrm{Tr}\left[m^{-1}(\mathbf{k})\right] = \frac{1}{d\,\hbar^2}\sum_{\alpha=1}^{d}\frac{\partial^2 E_n}{\partial k_\alpha^2}\,, \tag{17}$$

where d is the dimension; in this case, m^* reflects the average curvature in all directions and is equivalent to the effective mass used in the Drude model or Landau quantization. experimentally, m^* is directly accessible via cyclotron resonance measurements in magnetic fields: by tracking the resonance frequency

$$\omega_c = \frac{eB}{m^*}, \tag{18}$$

one extracts m^* from the slope of the Landau-level fan diagram.

Within the Boltzmann transport framework, the conductivity tensor at energy E takes the form

$$\sigma_{\alpha\beta}(E) = e^2\int_{\mathcal{B}} v_{n,\alpha}(\mathbf{k})\,v_{n,\beta}(\mathbf{k})\,\tau_n(\mathbf{k})\,\delta\big(E - E_n(\mathbf{k})\big)\,\frac{\mathrm{d}^d k}{(2\pi)^d}\,, \tag{19}$$

where $\tau_n(\mathbf{k})$ is the scattering time, directly linking band-structure curvature to measured mobilities and Hall measurements [1, 21]. This expression makes manifest how both the velocity $\mathbf{v}_n$ and the curvature $m^{-1}(\mathbf{k})$ underlie measured conductivities. For isotropic, parabolic bands one then recovers the familiar Drude-like behavior in terms of the scalar effective mass m^*:

$$\sigma = \frac{ne^2\tau}{m^*}, \tag{20}$$

DOI: 10.54254/2753-8818/2025.GL33588

However, in anisotropic or multi-valley systems, the full tensor structure $m_{\alpha\beta}$ is rendered crucial for the interpretation of Hall-effect measurements and mobility anisotropy. An intuitive real-space picture is provided by the velocity–dispersion relation itself: it is implied by a steeper slope in $E(\mathbf{k})$ that an applied electric field $\mathbf{E}$, which induces a shift $\hbar\dot{\mathbf{k}} = -e\mathbf{E}$, yields a larger rate of change in $\mathbf{r}$, corresponding to a higher drift velocity. Thus, device performance can be enhanced by band-structure engineering through strain, alloying, or heterostructuring, which optimizes $\nabla_{\mathbf{k}}E$. The empirical dispersion $E_n(\mathbf{k})$ and its slopes are directly mapped by modern angle-resolved photoemission spectroscopy (ARPES), from which both $\mathbf{v}_n$ and m^* are extracted across the Brillouin zone.

A bridge between microscopic band-structure theory and macroscopic transport observables is provided by the effective-mass tensor. The way in which lattice symmetry and electronic dispersion conspire to endow carriers with an apparent inertia is captured by it, and such inertia is directly measurable in cyclotron resonance, quantum oscillations, and Hall-effect experiments. Real-space dynamics under applied fields are likewise governed by the group velocity $\mathbf{v}_n(\mathbf{k})$, as the first derivative of $E(\mathbf{k})$, highlighting the central role of dispersion engineering in tailoring electronic transport.

2.4. Berry curvature and topological invariants

Beyond spectral and transport properties, the periodic parts $u_n(\mathbf{k})$ possess a geometric phase structure. Defining the Berry connection

$$\mathcal{A}_n(\mathbf{k}) = i\big\langle u_n(\mathbf{k}) \;\big|\; \nabla_{\mathbf{k}} u_n(\mathbf{k})\big\rangle, \tag{21}$$

one obtains the Berry curvature

$$\Omega_{n,\alpha\beta}(\mathbf{k}) = \partial_{k_\alpha}\mathcal{A}_{n,\beta} - \partial_{k_\beta}\mathcal{A}_{n,\alpha} = i\Big(\langle\partial_{k_\alpha}u_n|\partial_{k_\beta}u_n\rangle - \langle\partial_{k_\beta}u_n|\partial_{k_\alpha}u_n\rangle\Big). \tag{22}$$

Its integral over the Brillouin zone yields the Chern number

$$C_n = \frac{1}{2\pi}\int_{\mathcal{B}} \Omega_n(\mathbf{k})\,\mathrm{d}^2k, \tag{23}$$

quantized to integer values and signifying topological phases with protected edge states. In two-dimensional materials, by analytically continuing $u_{\mathbf{k}}$ into the complexified Brillouin zone, one obtains anomalous transverse responses, such as the spin Hall and quantum anomalous Hall effects at zero magnetic field [22, 23].

By recasting Bloch's theorem as a statement about one–dimensional unitary characters of the translation group, not only has the familiar form $\psi_{\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}}u_{\mathbf{k}}(\mathbf{r})$ been recovered, but a transparent algebraic underpinning for crystal momentum as an irreducible representation label has also been provided. Physically, it is highlighted by this perspective that band dispersion $E_n(\mathbf{k})$, group velocities $\mathbf{v}_n(\mathbf{k})$, and effective-mass tensors are emergent from the interplay between lattice symmetry and complex-analytic structure in reciprocal space. Methodologically, group-representation theory has been unified with analytic continuation techniques by our approach, paving the way for systematic lattice Green's-function computations via contour integrals in the complex k-plane. Moreover, by the inclusion of Berry-curvature and Chern-number concepts, this framework has been enriched through the revelation of geometric and topological aspects of Bloch bands. The Berry connection $\mathcal{A}_n(\mathbf{k}) = i\langle u_n(\mathbf{k})|\nabla_{\mathbf{k}}u_n(\mathbf{k})\rangle$ and its associated curvature $\Omega_{n,\alpha\beta}(\mathbf{k}) = \partial_{k_\alpha}\mathcal{A}_{n,\beta} - \partial_{k_\beta}\mathcal{A}_{n,\alpha}$ not only quantify anomalous transverse transport such as the spin Hall and quantum anomalous Hall effects at zero magnetic field, but are also the basis for quantized invariants that protect edge modes in

DOI: 10.54254/2753-8818/2025.GL33588

two-dimensional systems. By analytically continuing $u_{\mathbf{k}}$ into the complexified Brillouin zone, pole residues and branch-cut topology are further connected to topological circulation, thus blending residue-calculus methods with conformal-mapping approaches.

In sum, our algebraic–analytic synthesis deepens the understanding of Bloch states from first principles, equips the community with a versatile toolkit for spectral, transport, and topological calculations, and sets the stage for the residue and topology. A unified paradigm for tackling a broad class of problems in condensed-matter physics, from conventional band-structure analysis to the frontier of topological materials and beyond, is thus provided by the application of group representation theory in Bloch theory.

3. The topological perspective of the residue theorem and physical Interpretation

The classical residue theorem admits a succinct reinterpretation in terms of the topology and cohomology of the punctured complex plane. Let $D \subset \mathbb{C}$ be a connected domain and denote by

$$D' = D \setminus \{z_1, \ldots, z_n\} \tag{24}$$

the domain obtained by removing a finite collection of points. Each such singularity corresponds to an independent degree of freedom for looping, describing the topological properties of the loops around these singularities. These loops can naturally remind us of the homotopy group. From a homotopical perspective, D' is characterized by its fundamental group

$$\pi_1(D') \cong \mathbb{Z}^n, \tag{25}$$

generated by small, positively oriented loops γ_k around each removed point z_k. If C is any closed, piecewise smooth contour in D' whose homotopy class decomposes as

$$[C] = \sum_{k=1}^{n} I(C, z_k)[\gamma_k], \qquad I(C, z_k) \in \mathbb{Z}, \tag{26}$$

then the integral of a meromorphic one-form $\omega = f(z)\,dz$ depends solely on these winding numbers. Indeed, local analysis shows

$$\oint_{\gamma_k} f(z)\,dz = 2\pi i \operatorname{Res}(f, z_k), \tag{27}$$

and hence

$$\oint_C f(z)\,dz = 2\pi i \sum_{k=1}^{n} I(C, z_k) \operatorname{Res}(f, z_k). \tag{28}$$

Equivalently, in the language of de Rham cohomology [24, 25],

$$H^1_{\mathrm{dR}}(D') = \frac{\{\omega \in \Omega^1(D') : d\omega = 0\}}{\{d\phi : \phi \in C^\infty(D')\}}, \tag{29}$$

the cohomology class of ω is determined by its periods

$$\oint_{\gamma_k} \omega = 2\pi i \operatorname{Res}(f, z_k), \tag{30}$$

establishing a precise correspondence between analytic residues and topological invariants.

DOI: 10.54254/2753-8818/2025.GL33588

To illustrate the physical utility of this correspondence, we consider three canonical examples from electrostatics and fluid dynamics. In each case, the integral of a complex field around a closed contour yields a conserved quantity governed by the residue theorem.

First, we look the example about Electrostatic Field of a Point Charge. Consider a point charge q located at the origin in the two-dimensional plane. The potential $\phi(x, y)$ satisfies the 2D Poisson equation

$$\nabla^2\phi(x,y) \;=\; -\frac{q}{\varepsilon_{2D}}\,\delta(x)\,\delta(y), \tag{31}$$

and for $r = \sqrt{x^2+y^2} > 0$,

$$\nabla^2\phi = 0. \tag{32}$$

In polar coordinates (r, θ),

$$\nabla^2\phi = \frac{1}{r}\frac{\partial}{\partial r}\Big(r\,\frac{\partial\phi}{\partial r}\Big) + \frac{1}{r^2}\frac{\partial^2\phi}{\partial\theta^2}. \tag{33}$$

Assuming radial symmetry $\phi = \phi(r)$, the angular term vanishes and

$$\frac{1}{r}\frac{d}{dr}\Big(r\,\phi'(r)\Big) = 0 \quad\Longrightarrow\quad r\,\phi'(r) = A \quad\Longrightarrow\quad \phi'(r) = \frac{A}{r}. \tag{34}$$

Integrating gives

$$\phi(r) = A\ln r + B. \tag{35}$$

Matching the source via a contour integral,

$$\oint_{|\mathbf{r}|=R}\mathbf{E}\cdot d\boldsymbol{\ell} = -\oint_0^{2\pi}\phi'(R)\,R\,d\theta = -2\pi A = \frac{q}{\varepsilon_{2D}}, \tag{36}$$

we obtain

$$A = -\frac{q}{2\pi\,\varepsilon_{2D}}, \quad \phi(r) = -\frac{q}{2\pi\,\varepsilon_{2D}}\ln\Big(\frac{r}{r_0}\Big). \tag{37}$$

Since $\mathbf{E} = -\nabla\phi$,

$$\mathbf{E}(r) = \frac{q}{2\pi\,\varepsilon_{2D}}\frac{1}{r}\,\hat{\mathbf{r}} \quad\Longrightarrow\quad E_x = \frac{q}{2\pi\,\varepsilon_{2D}}\frac{x}{x^2+y^2}, \quad E_y = \frac{q}{2\pi\,\varepsilon_{2D}}\frac{y}{x^2+y^2}. \tag{38}$$

Now we can transform this real anlaysis representation to complex analysis form. Define the complex coordinate $z = x + iy$ and its conjugate $\bar{z} = x - iy$. Introduce the complex field

$$E(z) \;=\; E_x - iE_y \;=\; \frac{q}{2\pi\,\varepsilon_{2D}}\frac{x-iy}{x^2+y^2} \;=\; \frac{q}{2\pi\,\varepsilon_{2D}}\frac{1}{\bar{z}}. \tag{39}$$

Absorbing ε_{2D} into the constant and centering at z_0, we obtains the form

$$E(z) = \frac{q}{2\pi}\,\frac{1}{z-z_0} \quad (z_0 = 0), \tag{40}$$

We perform contour integration using the complex analysis form of the electric field.

$$\oint_C E(z)\,dz = 2\pi i\,I(C, z_0)\,\mathrm{Res}(E, z_0) = iqI(C, z_0) \tag{41}$$

when $I(C, z_0) = 1$ we get Gauss's law in the plane

DOI: 10.54254/2753-8818/2025.GL33588

Apart from the example of electromagnetism, we can also consider the example of fluid dynamics.Consider a single point vortex of strength Γ located at z_0 in an incompressible, inviscid two-dimensional flow. The complex potential for this vortex is

$$F(z) = -\frac{i\Gamma}{2\pi}\ln\big(z - z_0\big), \tag{42}$$

[26, 27] up to an additive constant. The complex velocity $w(z) = u - iv$ is given by

$$w(z) = \frac{dF}{dz} = -\frac{i\Gamma}{2\pi}\frac{1}{z - z_0} = \frac{i\Gamma}{2\pi}\frac{1}{z - z_0}, \tag{43}$$

with the sign convention for Γ indicating the sense of rotation. In Cartesian coordinates $z = x + iy$ and $z_0 = x_0 + iy_0$, one recovers the velocity components

$$u(x, y) = -\frac{\Gamma}{2\pi}\frac{y - y_0}{(x - x_0)^2 + (y - y_0)^2}, \quad v(x, y) = \frac{\Gamma}{2\pi}\frac{x - x_0}{(x - x_0)^2 + (y - y_0)^2}. \tag{44}$$

The circulation around any closed contour C enclosing the vortex is

$$\Gamma_C = \oint_C (u\,dx + v\,dy) = \Re\Big(\oint_C w(z)\,dz\Big) = \Re\big(2\pi i\,\mathrm{Res}(w, z_0)\big) = \Gamma\, I(C, z_0), \tag{45}$$

so when C winds once positively around z_0, $\Gamma_C = \Gamma$. This shows the circulation depends solely on the vortex strength and the winding number. We are now further considering the case of multiple vortices.For a system of n vortices at z_k with strengths Γ_k, the combined complex velocity field is

$$w(z) = \sum_{k=1}^{n} \frac{i\,\Gamma_k}{2\pi}\frac{1}{z - z_k}, \tag{46}$$

and the net circulation around C follows by linearity:

$$\oint_C w(z)\,dz = -\sum_{k=1}^{n} \Gamma_k\, I(C; z_k), \tag{47}$$

hence

$$\Gamma_C = \sum_{k=1}^{n} \Gamma_k\, I(C; z_k). \tag{48}$$

These examples underscore how local residues encode global, conserved physical observables.

In this section, the residue theorem is framed within the topological context of the fundamental group and de Rham cohomology, revealing how local singularity data of analytic functions are responsible for global invariants in physical systems. A departure from traditional approaches that rely on explicit path integrals or coordinate expansions is effected by this viewpoint; instead, the value of a contour integral is identified directly with the algebraic–topological combination of the residue at each singularity and the winding number of the contour. Consequently, when a contour is deformed within its homotopy class without crossing any singularity, the associated conserved quantity is preserved without the need for explicit reconstruction of integration paths. For the two-dimensional electrostatic problem, a point charge is represented by a simple pole of the complex electric field, its residue being proportional to the charge. The contour integral of the complex field around any closed

DOI: 10.54254/2753-8818/2025.GL33588

loop is shown to reproduce the planar Gaussian law, establishing that the electric flux enclosed by the loop is proportional to the enclosed charge times the winding number. When the loop does not encircle the singularity, the net flux is found to vanish regardless of the loop shape. This complex-analytic derivation not only recovers the standard divergence-based proof of Gauss's theorem but also provides an explanation, via homotopy invariance, of why the flux is independent of the contour's detailed shape. Compared to the method that invokes the divergence theorem and area-element integration, the residue-based perspective illustrates the internal consistency from singularity to residue and then to global flux, and it is naturally extended to configurations with multiple or distributed singularities: it is only required that each pole's residue and its winding number be identified, thereby avoiding repeated differential-element integrations.

In fluid dynamics, the complex potential of a single point vortex is expressed by a logarithmic function, whose derivative produces a simple pole in the complex velocity field with the residue equal to the vortex strength. Upon application of the residue theorem to a closed contour, Kelvin's circulation theorem is obtained: provided that the contour is deformed without crossing the vortex location, the circulation is fixed at the product of vortex strength and winding number. Explicit coordinate-by-coordinate integration or construction of streamlines is not required. When the formulation is generalized to multiple vortices, the complex velocity is represented as the sum of several simple poles, and the circulation around any contour is given by the sum of the individual residues weighted by their respective winding numbers. This topological–cohomological method provides a concise treatment of global circulation in interacting vortex systems and facilitates the analysis of how circulation invariants evolve when vortices move or contours deform: changes in homotopy class directly indicate possible jumps or reconnections in circulation, thereby shedding light on the flow's stability and topological evolution. Compared with traditional approaches that often require solving for the velocity potential in each specific geometry or boundary condition followed by explicit integrals, the residue-based framework requires only knowledge of vortex positions, strengths, and contour winding; this is especially advantageous when boundaries are complex or singularities are numerous. Moreover, this approach can be used to inform numerical simulations by first identifying singularity locations and their topological arrangement, and then imposing global circulation constraints via residue computations to improve efficiency and stability. Viewed more broadly, the unification of residue calculus and homotopy invariance in the explanation of conserved quantities is seen to resonate with themes in modern mathematical physics, such as conformal field theory and topological phases of matter. For example, Berry phases and geometric phases are often interpreted through manifold cohomology or fiber-bundle structures; similarly, the invariance of vortex circulation or electric flux under continuous deformation is paralleled by the role of topological invariants such as Chern numbers in quantum Hall systems. The emphasis of the present method on "local singularity properties leading to global topological invariants" may inform studies of quantum vortices or topologically protected states in many-body systems. In nonlinear or dissipative contexts, if singularity evolution corresponds to branch points or singular manifolds, cohomological techniques could be employed to identify mechanisms of conservation breaking or topological transitions. Thus, the algebraic–topological framework developed here is applicable not only to classical electrostatics and ideal fluid flows but also lays the conceptual groundwork for more complex settings, such as lattice Green's function problems on branched Riemann surfaces, electromagnetic field distributions in nontrivial geometries, and field theories on higher-dimensional complex manifolds.

In summary, the derivation based on residue calculus and topological homotopy invariance not only reproduces and deepens the understanding of key conservation laws in electrostatics and fluid dynamics but also is seen to offer clear advantages in conceptual clarity, extensibility, and the handling

DOI: 10.54254/2753-8818/2025.GL33588

of complex singularity structures. By reducing conserved quantities to a unified combination of local algebraic data and global topological information, a succinct yet powerful theoretical toolset for the exploration of singularities and conservation properties across a wide range of physical problems is provided by this approach.

4. Conformal mapping and its applications in electrodynamics

4.1. Mathematical background of conformal mapping and the electrostatic wedge problem

In addition to the aforementioned complex analysis content, the part of conformal mapping in complex analysis also plays a significant role in the field of physics. We will now discuss this part.We begin with the standard definition of a conformal mapping in the plane. Let $z = x + iy$ denote the complex coordinate, and let $w = f(z)$ be a holomorphic function with nonvanishing derivative:$f'(z) \neq 0$ in the region of interest Writing

$$f(z) = u(x, y) + iv(x, y), \tag{49}$$

the Cauchy–Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}, \tag{50}$$

ensure that f is holomorphic. The Jacobian matrix of the mapping is

$$J_f(z) = \begin{pmatrix} \Re f'(z) & -\Im f'(z) \\ \Im f'(z) & \Re f'(z) \end{pmatrix}, \tag{51}$$

so that in the tangent plane df acts locally as a rotation by $\arg f'(z)$ followed by a uniform dilation by $|f'(z)|$. Consequently, conformal mappings preserve all intersection angles between smooth curves (including orthogonality), making them powerful tools for transforming boundary value problems in two dimensions. Next, we will demonstrate the advantages of the conformal mapping method by using a static problem from electrodynamics.Consider the two-dimensional electrostatic potential $\Phi(r, \theta)$ in a wedge-shaped region:$\mathcal{W} = \{(r, \theta) \mid 0 < r < \infty,\ -\alpha \leq \theta \leq \alpha\}$, in polar coordinates (r, θ). The potential satisfies Laplace's equation,

$$\nabla^2 \Phi = \frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial \Phi}{\partial r}\right) + \frac{1}{r^2}\frac{\partial^2 \Phi}{\partial \theta^2} = 0, \tag{52}$$

subject to Dirichlet boundary conditions on the conducting faces:

$$\Phi(r, \theta = \pm\alpha) = 0 \quad (\forall r > 0). \tag{53}$$

In order to simplify the geometric boundary conditions in the problem, we employed the conformal mapping method.First we Introduce the complex variable

$$\zeta = re^{i\theta}, \tag{54}$$

so that the wedge $\mathcal{W}$ in the ζ-plane corresponds to the angular sector $-\alpha \leq \arg\zeta \leq \alpha$. Define the conformal map

$$w = \zeta^{\frac{\pi}{2\alpha}} = r^{\frac{\pi}{2\alpha}} e^{i\frac{\pi}{2\alpha}\theta}. \tag{55}$$

Under this map the boundaries $\theta = \pm\alpha$ are sent to the real axis in the w-plane, since $\arg w = \frac{\pi}{2\alpha}(\pm\alpha) = \pm\frac{\pi}{2}$, but in fact the images lie on the real axis up to a multiplication by $e^{\pm i\pi/2}$, which can

DOI: 10.54254/2753-8818/2025.GL33588

be rotated back to the real axis by a constant phase; equivalently, one may compose with a rotation of the w-plane if desired. The interior $-\alpha < \theta < \alpha$ is mapped onto the upper half-plane $\Im(w) > 0$. Since conformal maps preserve harmonic functions which is the solutions of Laplace's equation up to composition, the potential $\Phi(r, \theta)$ in the wedge corresponds to a harmonic function $\Psi(w)$ in the upper half-plane satisfying

$$\frac{\partial^2 \Psi}{\partial w \partial \bar{w}} = 0, \quad \Psi(w) = 0 \quad \text{for } \Im(w) = 0. \tag{56}$$

Next we can get move to the solution. A general harmonic function in the upper half-plane vanishing on the real axis can be expressed in terms of the argument of w. A standard choice is

$$\Psi(w) = A \arg(w), \quad \arg(w) \in (0, \pi), \tag{57}$$

where A is a constant to be fixed by the prescribed potential scale. Pulling back to the ζ-plane via $w = \zeta^{\pi/(2\alpha)}$, we write

$$\Psi(w) = A\Im\big(\ln w\big) = A\Im\Big(\ln\big(r^{\frac{\pi}{2\alpha}} e^{i\frac{\pi}{2\alpha}\theta}\big)\Big) = A\frac{\pi}{2\alpha}\theta. \tag{58}$$

Hence the original potential in the wedge is

$$\Phi(r, \theta) = A\,\frac{\pi}{2\alpha}\theta. \tag{59}$$

If we require $\Phi(r, \theta = \alpha) = \Phi_0$ on the boundary $\theta = +\alpha$, then

$$A\frac{\pi}{2\alpha} \cdot \alpha = \Phi_0 \implies A = \frac{2\alpha}{\pi}\,\Phi_0.$$

Thus

$$\Phi(r, \theta) = \Phi_0\,\frac{\theta}{\alpha}, \quad -\alpha \le \theta \le \alpha. \tag{60}$$

We observe that Φ is independent of r and depends linearly on θ. And electric field is $\mathbf{E} = -\nabla\Phi$. In polar coordinates,

$$E_r = -\frac{\partial \Phi}{\partial r} = 0, \qquad E_\theta = -\frac{1}{r}\frac{\partial \Phi}{\partial \theta} = -\frac{1}{\alpha r}. \tag{61}$$

Thus the magnitude $|\mathbf{E}| = \frac{1}{\alpha r}$,which diverges as $r \to 0$, indicating a singular field enhancement near the wedge tip. The sharper the wedge , the stronger the enhancement for a given small r.

The charge density, as an important physical quantity, can also be obtained from the above solution. On each conducting face $\theta = \pm\alpha$, the surface charge density σ is given by

$$\sigma = \epsilon_0\,\frac{\partial \Phi}{\partial n}. \tag{62}$$

Since $\Phi = \theta/\alpha$, along $\theta = \pm\alpha$ the outward normal derivative (pointing into the region) has magnitude $\partial\Phi/\partial n = 1/\alpha$. Therefore,

$$\sigma = \epsilon_0\,\frac{1}{\alpha}. \tag{63}$$

This boundary charge produces the field in the wedge. As $\alpha \to 0$, the surface charge density increases, consistent with the field enhancement near the tip.

The solution $\Phi(r, \theta) = (\theta/\alpha)\Phi_0$ is noted for its simplicity: independence of r and linearity in θ are exhibited. Physically, it is indicated that the potential in the wedge is entirely dictated by the boundary

DOI: 10.54254/2753-8818/2025.GL33588

values on the conducting faces, with no volumetric charge in the interior. The divergence of $|\mathbf{E}|$ as $r \to 0$ is reflective of the classical field enhancement at sharp tips, which has been shown to have important implications in microelectronics, nanostructures , and vacuum electronics. The explicit form demonstrates how the wedge angle α controls the strength of the singularity: larger fields near the apex are produced when α is smaller. From a methodological standpoint, the Laplace equation in a complicated domain is transformed by means of conformal mapping into an equivalent problem in a canonical domain where standard solutions are known. Compared to traditional separation-of-variables approaches applied directly in the wedge, which also yield a Fourier-type series or an ansatz $\Phi \sim r^n \sin(n\theta/\alpha)$, the conformal mapping approach is often found to yield the simplest derivation and clearer geometric insight: the mapping of the angular sector to a half-plane is directly visualized and the harmonic solution is supplied by the argument function. In more intricate geometries, for example in domains with multiple corners, curved boundaries, or mixed boundary conditions, the problem can be rendered tractable by conformal mapping to regions where boundary conditions are transformed into straight lines or circles, thereby enabling the employment of known Green's functions or Poisson integrals. Thus, both an elegant derivation and a conceptual framework highlighting the role of geometry in the potential distribution are provided by conformal mapping.

4.2. Isomorphism between the Möbius Transformation group and the lorentz group

Conformal mapping is not merely used to solve partial differential equation problems; it can also provide us with an enlightening understanding of the algebraic structure of physical laws from the perspective of complex analysis. One example is that it reveals the form of the Lorentz transformation determined by the spacetime symmetry in Minkowski spacetime from the perspective of complex analysis, which is of great significance for the development of quantum electrodynamics.

We first introduce Möbius transformations and $\mathrm{PSL}(2,\mathbb{C})$ A Möbius transformation is a map on the extended complex plane $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ of the form

$$z \;\mapsto\; w = \frac{az+b}{cz+d}, \quad a,b,c,d \in \mathbb{C}, \quad ad-bc \neq 0. \tag{64}$$

Such transformations form a group under composition, and by identifying $\lambda(a,b,c,d)$ with (a,b,c,d) for nonzero $\lambda \in \mathbb{C}^\times$, one obtains the projective special linear group $\mathrm{PSL}(2,\mathbb{C}) \cong \mathrm{SL}(2,\mathbb{C})/\{\pm I\}$. These maps act conformally on $\widehat{\mathbb{C}}$, mapping circles or lines to circles or lines, preserving cross-ratios, and so on. Next we introduce Lorentz transformations via $\mathrm{SL}(2,\mathbb{C})$ In four-dimensional Minkowski space with signature $(+,-,-,-)$, a real four-vector x^μ may be represented by a Hermitian 2×2 matrix:

$$X = x^0\sigma_0 + x^1\sigma_1 + x^2\sigma_2 + x^3\sigma_3 = \begin{pmatrix} x^0 + x^3 & x^1 - ix^2 \\ x^1 + ix^2 & x^0 - x^3 \end{pmatrix}, \tag{65}$$

where $\sigma_0 = I$ is the identity and σ_i are the Pauli matrices. The determinant

$$\det X = (x^0)^2 - (x^1)^2 - (x^2)^2 - (x^3)^2 \tag{66}$$

is the Minkowski norm. For any $A \in \mathrm{SL}(2,\mathbb{C})$, define

$$X \;\mapsto\; X' = A\,X\,A^\dagger. \tag{67}$$

Since $\det X' = \det A \, \det X \, \det A^\dagger = \det X$ and X' remains Hermitian, this action corresponds to a proper orthochronous Lorentz transformation on the four-vector x^μ. The mapping $A \mapsto \Lambda(A) \in \mathrm{SO}^+(1,3)$ has kernel $\{\pm I\}$, yielding the exact sequence

$$1 \;\longrightarrow\; \{\pm I\} \;\longrightarrow\; \mathrm{SL}(2,\mathbb{C}) \;\longrightarrow\; \mathrm{SO}^+(1,3) \;\longrightarrow\; 1, \tag{68}$$

DOI: 10.54254/2753-8818/2025.GL33588

and hence

$$\mathrm{SO}^+(1,3) \;\cong\; \mathrm{SL}(2,\mathbb{C})/\{\pm I\} \;\cong\; \mathrm{PSL}(2,\mathbb{C}) \;\cong\; \mathrm{M\ddot{o}b}(\widehat{\mathbb{C}}). \tag{69}$$

The identification $\mathrm{PSL}(2,\mathbb{C}) \cong \mathrm{SO}^+(1,3)$ has long been recognized as a deep and classical result that reveals a profound interplay between two-dimensional complex analysis and four-dimensional Minkowski geometry. On one hand, the Riemann sphere is acted upon conformally by Möbius transformations, with angles and circles being preserved; on the other hand, the light-cone structure in spacetime is preserved by Lorentz transformations. Mathematically, it is implied by this isomorphism that any proper orthochronous Lorentz transformation can be realized as the conjugation action of some $A \in \mathrm{SL}(2,\mathbb{C})$ on Hermitian matrices representing four-vectors. Topologically, $\mathrm{SL}(2,\mathbb{C})$ is presented as a double cover of $\mathrm{SO}^+(1,3)$, which explains the two-to-one correspondence $\pm A \mapsto \Lambda$. In representation theory, the spinor structure in relativistic theories is underlied by this correspondence: spin-$\frac{1}{2}$ representations are realized in the fundamental of $\mathrm{SL}(2,\mathbb{C})$, while integer-spin representations are descended to representations of $\mathrm{SO}^+(1,3)$. This result is taken to set the stage for physical applications in which massless particles and conformal structures are interplayed. In particular, null directions in Minkowski space are identified with points on the Riemann sphere via stereographic projection; Möbius transformations are then employed to describe the action of Lorentz transformations on this sphere, leading to insights in scattering amplitudes and in the celestial sphere approach to quantum field theory.

Now, let's use the physical example we just discussed to demonstrate this isomorphism relationship. We work in natural units ($c = 1$) and consider a future-directed, null four-momentum for a photon:

$$P^\mu = \omega\,(1,\,\hat{n}), \quad \hat{n} = (\sin\theta\cos\phi,\ \sin\theta\sin\phi,\ \cos\theta), \quad \omega > 0. \tag{70}$$

Each direction $\hat{n}$ on the celestial sphere can be labelled by stereographic coordinates z on the Riemann sphere:

$$z = e^{i\phi}\tan\frac{\theta}{2}. \tag{71}$$

Here $z \in \mathbb{C}$ covers all directions except the "south pole" ($\theta = \pi$), with $z \to \infty$ corresponding to that point. Thus there is a one-to-one correspondence between null directions of photon momenta (up to overall scaling ω) and points $z \in \widehat{\mathbb{C}}$.

According to our purpose let us discuss Lorentz action via $\mathrm{SL}(2,\mathbb{C})$ and Möbius transformations. Recall the Hermitian matrix representation of four-vectors:

$$P = P^\mu \sigma_\mu, \quad \sigma_\mu = (\mathbf{1}, \sigma_i). \tag{72}$$

For null P^μ, $\det P = 0$, so P factors as

$$P = \omega\,\xi(z)\,\xi^\dagger(z), \quad \xi(z) = \begin{pmatrix} 1 \\ z \end{pmatrix}. \tag{73}$$

An overall complex phase of $\xi(z)$ does not change P but corresponds to the $U(1)$ freedom of the spinor. Under $A \in \mathrm{SL}(2,\mathbb{C})$, we have

$$P \;\mapsto\; P' = A\,P\,A^\dagger = \omega\,(A\xi(z))\,(A\xi(z))^\dagger.$$

Since $A\xi(z)$ is proportional to another spinor $\xi(z')$, one reads off the transformed stereographic coordinate:

$$z' \;=\; \frac{az+b}{cz+d}, \quad A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{C}). \tag{74}$$

DOI: 10.54254/2753-8818/2025.GL33588

Thus Lorentz transformations act on null directions exactly as Möbius transformations on $\widehat{\mathbb{C}}$. A plane-wave solution for a massless scalar or photon field may be written as

$$\Phi_P(x) = e^{-iP\cdot x}. \tag{75}$$

Under a Lorentz transformation corresponding to $z \mapsto z'$, the coordinate x transforms, and one finds that $\Phi_P(x)$ transforms up to a phase and a local dilation factor related to the Jacobian of the Möbius map. In celestial conformal field theory language, this corresponds to the field having conformal weight -1 on the sphere, reflecting how amplitudes transform under the conformal action on the celestial sphere.

The derivation above concretely demonstrates that null directions (photon trajectories) are transformed under Lorentz transformations exactly as points on the Riemann sphere are transformed under Möbius maps [28, 29]. This phenomenon is seen to directly reflect the previously established isomorphism $\mathrm{PSL}(2,\mathbb{C}) \cong \mathrm{SO}^+(1,3)$. Physically, it is implied that for massless particles, the space of asymptotic directions at null infinity is naturally a conformal sphere, and that Lorentz symmetry is realized as the conformal symmetry on that sphere. This insight is taken to underpin modern developments in scattering amplitudes and the so-called "celestial sphere" program, in which four-dimensional scattering processes are studied by mapping them onto correlation functions of a two-dimensional conformal field theory living on the celestial sphere at null infinity. In this picture: First, an insertion at a point z on the sphere is associated with each external massless particle, with conformal weights being determined by the particle's energy and spin. Second, the insertion points are transformed conformally by Lorentz transformations, constraining the functional form of amplitudes. Third, extensions to include asymptotic symmetries such as the BMS group are made, enriching this structure and relating soft theorems in four dimensions to Ward identities in the two-dimensional theory. Thus, concrete implications are possessed by the Möbius–Lorentz correspondence: the transformation of physical data at infinity is dictated by it, the decomposition of amplitudes into conformal correlators is determined by it, and the manifestation of symmetry constraints in the celestial CFT framework is prescribed by it. Furthermore, the conformal invariance of massless field equations under the action at null infinity is clarified by this viewpoint. It is also suggested that geometric structures from complex analysis find direct application in describing scattering processes, encoding the dependence on kinematic invariants in a manner that is manifestly covariant under Lorentz symmetry.

5. Conclusion

In this work, algebraic, topological, and analytic strands of complex analysis and group theory have been woven together to offer a unified treatment of three prototypical problems in mathematical physics. First, Bloch's theorem was cast as a statement about one–dimensional unitary characters of the translation group, thereby illuminating how crystal momentum is emerged as an irreducible representation label and how spectral, transport, and topological properties of electronic bands are directly informed. Next, the residue theorem was reframed within the language of homotopy and de Rham cohomology, showing that conserved quantities such as planar electric flux and vortex circulation are encoded purely by local residues and winding numbers, independent of the contour's shape. Finally, via a judicious conformal map, the conducting-wedge electrostatic problem was solved in closed form, the characteristic $1/r$ singularity at the apex was elucidated, and the deep isomorphism between Möbius transformations and the proper orthochronous Lorentz group was unveiled, providing the mathematical backbone for modern celestial sphere approaches to massless scattering amplitudes.

Looking ahead, two concrete avenues for further exploration are identified. First, the residue–homotopy framework can be extended to nonlinear and interacting field theories: by combining

DOI: 10.54254/2753-8818/2025.GL33588

contour-integral methods with Riemann-surface techniques, branch-point collisions and topological transitions in many-body systems may be tracked. Second, generalization to three-dimensional and quasi-conformal settings is invited by the conformal-mapping and Möbius–Lorentz correspondence, enabling analytic and numerical advances in complex boundary-value problems such as electrostatic metamaterials or gravitational lensing, and furnishing higher-dimensional spinor analogues for use in curved spacetime. It is anticipated that these targeted developments will continue to bridge abstract complex-analytic invariants with tangible physical phenomena, advancing both theory and applications across condensed-matter physics, fluid dynamics, and quantum field theory.

Acknowledgments

I would like to express my gratitude to Dr.Chenran Gu for her guidance and assistance in the writing of this article. Her academic advice has greatly inspired me.

References

[1] N. W. Ashcroft and N. D. Mermin (1976), *Solid State Physics*. Saunders College Publishing, Philadelphia.
[2] M. S. Dresselhaus, G. Dresselhaus, and A. Jorio (2008), *Group Theory: Application to the Physics of Condensed Matter*. Springer.
[3] J. D. Jackson (1999), *Classical Electrodynamics*, 3rd ed. Wiley.
[4] E. M. Stein and R. Shakarchi (2003), *Complex Analysis*. Princeton University Press.
[5] F. Bloch (1929), "Über die Quantenmechanik der Elektronen in Kristallgittern," *Zeitschrift für Physik*, **52**(7-8): 555–600.
[6] J. Zak (1964), "Magnetic Translation Group," *Physical Review*, **134**(6A): A1602–A1611.
[7] C. Kittel (2004), *Introduction to Solid State Physics*, 8th ed. Wiley.
[8] N. I. Muskhelishvili (1992), *Singular Integral Equations: Boundary Problems of Function Theory and Their Application to Mathematical Physics*. Dover Publications.
[9] P. J. Olver (1995), *Equivalence, Invariants, and Symmetry*. Cambridge University Press.
[10] Z. Nehari (1952), *Conformal Mapping*. Dover Publications.
[11] G. K. Batchelor (2000), *An Introduction to Fluid Dynamics*. Cambridge University Press.
[12] G. B. Arfken and H. J. Weber (2005), *Mathematical Methods for Physicists*, 6th ed. Elsevier.
[13] P. M. Morse and H. Feshbach (1953), *Methods of Theoretical Physics*, Parts I & II. McGraw-Hill Book Company, New York.
[14] L. Ahlfors (1979), *Complex Analysis*, 3rd ed. McGraw-Hill.
[15] J. B. Conway (1978), *Functions of One Complex Variable*, 1st ed. Springer.
[16] W. Rudin (1987), *Real and Complex Analysis*, 3rd ed. McGraw-Hill.
[17] M. Hamermesh (1989), *Group Theory and Its Application to Physical Problems*. Dover Publications.
[18] H. Lamb (1932), *Hydrodynamics*, 6th ed. Cambridge University Press.
[19] E. N. Economou (2006), *Green's Functions in Quantum Physics*, 3rd ed. Springer.
[20] G. Grosso and G. P. Parravicini (2000), *Solid State Physics*. Academic Press.
[21] J. M. Ziman (1972), *Principles of the Theory of Solids*. Cambridge University Press.
[22] M. V. Berry (1984), "Quantal Phase Factors Accompanying Adiabatic Changes," *Proceedings of the Royal Society of London A*, **392**(1802): 45–57.
[23] D. Xiao, M.-C. Chang, and Q. Niu (2010), "Berry Phase Effects on Electronic Properties," *Reviews of Modern Physics*, **82**(3): 1959–2007.
[24] R. Remmert (1991), *Theory of Complex Functions*. Springer.
[25] R. Bott and L. W. Tu (1982), *Differential Forms in Algebraic Topology*. Springer.
[26] J. W. Brown and R. V. Churchill (2009), *Complex Variables and Applications*, 8th ed. McGraw-Hill.
[27] P. G. Saffman (1992), *Vortex Dynamics*. Cambridge University Press.
[28] H. M. Farkas and I. Kra (1992), *Riemann Surfaces*, 2nd ed. Springer.
[29] J. S. Walker (2004), *Complex Variables with Applications*, 5th ed. Pearson.