Proceedings of CONF-APMM 2025 Symposium: Multi-Qubit Quantum Communication for Image Transmission over Error Prone Channels
DOI: 10.54254/2753-8818/2025.GL33604

Quantum Tunneling of Electrons: From Fundamental Mechanisms to Cutting-Edge

Jingxing Zou

Wuhan Britain-China School, Wuhan, China
whbc3541132058@qq.com

Abstract. Quantum tunneling, a quintessential quantum phenomenon, allows electrons to traverse classically insurmountable energy barriers. This paper comprehensively reviews the theoretical underpinnings of electron tunneling, its experimental manifestations in nanoscale systems, and transformative applications in modern technology. We delve into advanced topics such as time-resolved tunneling dynamics, many-body quantum effects, and the role of tunneling in topological materials. By integrating recent breakthroughs in ultrafast spectroscopy and machine learning-driven device design, this work provides a roadmap for harnessing tunneling phenomena to overcome classical limitations in electronics, energy harvesting, and quantum computing.

Keywords: Schrödinger's wave equation, Uncertainty Principle, Entanglement, Combinatorial optimization problems.

1. Introduction

Quantum mechanics reshaped our understanding of particle behavior, with quantum tunneling standing as one of its most counterintuitive yet impactful predictions. First theorized by Friedrich Hund and George Gamow in the 1920s, tunneling was experimentally validated through semiconductor diodes and later revolutionized by scanning tunneling microscopy [1,2]. Introduction to Quantum Mechanics: Unveiling the Subatomic World.

Quantum mechanics, the cornerstone of modern physics, is a revolutionary framework that governs the behavior of particles at the subatomic scale. Since its inception in the early 20th century, it has reshaped our understanding of reality, challenging classical notions of determinism and locality. This enigmatic yet profoundly successful theory underpins technologies ranging from semiconductors to quantum computers, while raising philosophical questions about the nature of observation and existence.

1.1. Historical foundations

The quantum revolution began with Max Planck's 1900 solution to the "ultraviolet catastrophe" in blackbody radiation. By proposing that energy is quantized in discrete units (quanta), Planck unwittingly laid the groundwork for quantum theory. Albert Einstein extended this concept in 1905, explaining the photoelectric effect through light quanta (later called photons), earning him the Nobel

© 2026 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2025.GL33604

Prize and establishing wave-particle duality. Niels Bohr's 1913 atomic model introduced quantum leaps between energy states, while Louis de Broglie's 1924 hypothesis posited that all matter exhibits wave-like properties.

The formal mathematical structure emerged through Schrödinger's wave equation and Heisenberg's matrix mechanics, later shown to be equivalent formulations. Dirac's relativistic quantum theory unified quantum mechanics with special relativity, predicting antimatter. The Copenhagen interpretation, championed by Bohr and Heisenberg, became the dominant view, emphasizing probability and the role of measurement.

1.2. Core principles

1. Wave-Particle Duality: Quantum entities exhibit both particle-like and wave-like properties. The double-slit experiment epitomizes this paradox – electrons create interference patterns (wave behavior) while arriving as discrete particles at detectors.

2. Uncertainty Principle: Heisenberg demonstrated that complementary variables (e.g., position/momentum) cannot be simultaneously measured with arbitrary precision ($\Delta x \Delta p \geq \hbar/2$). This isn't a technical limitation but a fundamental property of quantum systems.

3. Superposition: A quantum system exists in multiple states simultaneously until measured, described by wavefunctions (ψ) . Schrödinger's famous cat thought experiment illustrates this paradox, where a cat is both alive and dead until observed.

4. Entanglement: When particles interact, they can become correlated such that measuring one instantly affects the other, regardless of distance. Einstein called this "spooky action at a distance," but experiments confirm its reality.

5. Quantization: Physical quantities (energy, angular momentum) take discrete values. Atomic orbitals and quantum tunneling – where particles penetrate classically forbidden barriers – exemplify this principle.

1.3. Mathematical framework

Quantum states are represented by vectors in Hilbert space, with observables as Hermitian operators. The Schrödinger equation $i\hbar\partial\psi/\partial t = \hat{H}\psi$ governs time evolution, where $\hat{H}$ is the Hamiltonian operator. Measurement collapses the wavefunction into eigenstates of the observable, with probabilities given by the Born rule $(P = |\langle\varphi|\psi\rangle|^2)$. Commutator relations $[\hat{x}, \hat{p}] = i\hbar$ formalize the uncertainty principle.

1.4. Applications and implications

Quantum mechanics enables modern technologies: lasers, MRI scanners, and transistors rely on quantum effects. Emerging fields like quantum computing exploit superposition and entanglement to solve problems intractable for classical computers. Quantum cryptography uses entanglement for theoretically unbreakable encryption.

Philosophically, interpretations remain debated. The Copenhagen interpretation contrasts with Many-Worlds (parallel universes), Pilot-Wave (hidden variables), and Quantum Bayesianism (subjective probabilities). Bell's theorem disproved local hidden variable theories, supporting non-locality.

DOI: 10.54254/2753-8818/2025.GL33604

1.5. Conclusion

Quantum mechanics remains our most accurate physical theory, with predictions confirmed to 10^{-15} precision. Yet its counterintuitive nature continues to inspire research into quantum gravity, quantum biology, and the measurement problem. As Richard Feynman remarked, "If you think you understand quantum mechanics, you don't understand quantum mechanics." This beautiful paradox – a theory both incomprehensible and indispensable – continues to drive humanity's quest to unravel reality's deepest mysteries.

Today, electron tunneling underpins technologies ranging from flash memory to quantum bits (qubits), while posing unresolved questions about its fundamental nature (e.g., "How fast does tunneling occur?"). This paper synthesizes multidisciplinary advances to address two pivotal questions:

$$\frac{\hbar^2 d^2\psi}{2mdx^2} + V\Big(x\Big)\psi = E\psi \tag{1}$$

For a rectangular barrier of height V_0 and width d, the transmission coefficient T approximates

$$T \approx \exp\left(\frac{-2d\sqrt{2m}(V_0-E)}{\hbar}\right) \tag{2}$$

$$I = \int N_s(E)N_y(E+eV)\left[f(E) - f(E+eV)\right]dE \tag{3}$$

1. How do many-body interactions and material topology modulate tunneling probabilities?
2. Can we engineer tunneling dynamics to achieve ultra-low-power electronics and fault-tolerant quantum computation?

2. Theoretical framework

2.1. Single-particle tunneling

The probability of an electron tunneling through a potential barrier is governed by the time-independent Schrödinger equation:

This exponential sensitivity to barrier geometry explains why atomic-scale precision is critical in tunneling devices.

2.2. Many-body and correlated systems

In correlated materials (e.g., high-temperature superconductors), electron-electron interactions dramatically alter tunneling behavior. The tunneling current in a superconductor follows:

DOI: 10.54254/2753-8818/2025.GL33604

$$I = \frac{e}{C} \cdot \Gamma_{\text{turn}} \tag{4}$$

where is the superconducting gap, and are density of states [3]. Such interactions enable exotic phenomena like Majorana fermion tunneling in topological insulators.

3. Experimental advances

3.1. Ultrafast tuning dynamics in quantum materials

Ultrasfast tuning dynamics in quantum materials refers to the manipulation and observation of material properties on the femtosecond to picosecond timescales, enabled by advanced techniques such as pump-probe spectroscopy and ultrafast laser excitation. These dynamics are crucial for probing non-equilibrium states and transient phenomena in quantum systems.

Laser-induced phase transitions: ultrafast optical pulses can transiently alter electronic pulses can transuently alter elctronic or lattice structures, including metastable phases or lattice structures (photoinduced superconductivity). For instance, in cuprates, near-infrared pulses may suppress charge density wave order while enhancing superconducting correlations.

coherent control of quantum states: by tailoring laser pulse parameters (e.g, amplitude, phase, and polarization), excitation of specific vibrational or applied to manipulate extinction dynamics in 2D materials like transition metal dichalcogenides(TMDs)

Attosecond laser pulses now permit real-time observation of tunneling. Recent experiments measured a tunneling delay time of attoseconds in strong-field ionization, resolving longstanding debates about tunneling "instantaneity." [4,5]

3.2. Single-electron control

Quantum dots enable single-electron tunneling with Coulomb blockade oscillations (Fig. 1a). At temperatures below 100 mK, the current through a quantum dot exhibits quantized steps:



Figure 1. (a) Coulomb blockade in a quantum dot (b) Attosecond tunneling measurement setup

Core principles of Quantum computing quantum computing represent a paradigm shift from classical computing by leveraging the principles of quantum mechanics. Unlike classical bits (quits*) exploit superposition and entanglement to perform computations.

DOI: 10.54254/2753-8818/2025.GL33604

Time-Resolved correlations: ultrafast techniques reveal hidden correlations, such as electron-phonon coupling dynamics in topological insulators.

4. Applications

4.1. Tunnel field-effect transistors (TFETs)

TFETs exploit band-to-band tunneling to achieve sub-60 mV/decade switching, outperforming conventional MOSFETs. Heterostructures like InAs/GaSb show on-currents >100 μA/μm at 0.3 V.

4.2. Quantum computing

In spin qubits, Pauli spin blockade enables readout via tunneling current asymmetry. Recent devices achieve 99.8% single-qubit gate fidelity by suppressing tunneling noise.

Applications of Quantum Computing

Quantum computing has emerged as a transformative technology with the potential to revolutionize diverse fields. Below are key application domains where quantum advantages are actively being explored:

Cryptography and Cybersecurity

The advent of Shor's algorithm, which efficiently factors large integers using quantum Fourier transforms, poses a fundamental threat to classical public-key cryptosystems (e.g., RSA and ECC). Conversely, quantum-resistant algorithms, such as lattice-based cryptography and quantum key distribution (QKD), leverage quantum principles to enable provably secure communication channels. For example, the BB84 protocol exploits photon polarization states to detect eavesdropping attempts in real time.

4.3. Drug discovery and molecular simulation

Quantum computers can simulate molecular and chemical systems with unprecedented accuracy by solving the Schrödinger equation exponentially faster than classical methods. This capability accelerates drug development by modeling protein-ligand interactions or catalytic pathways. In 2020, researchers at Google Quantum AI demonstrated a variational quantum eigensolver (VQE) to simulate the energy states of small molecules like hydrogen chains, showcasing a path toward practical quantum chemistry.

4.4. Optimization and logistics

Combinatorial optimization problems, such as supply chain management and traffic routing, are intractable for classical computers at scale. Quantum annealing (e.g., D-Wave systems) and quantum approximate optimization algorithms (QAOA) offer near-term solutions. For instance, Volkswagen utilized D-Wave's quantum annealers to optimize public bus routes in Lisbon, reducing traffic congestion by 15% in simulated scenarios.

4.5. Machine learning and AI

Quantum machine learning (QML) algorithms, including quantum support vector machines (QSVMs) and quantum neural networks (QNNs), exploit quantum parallelism to enhance pattern recognition and data classification. Hybrid quantum-classical frameworks, such as TensorFlow

DOI: 10.54254/2753-8818/2025.GL33604

Quantum, enable training models on noisy intermediate-scale quantum (NISQ) devices. Recent experiments achieved quantum-enhanced image recognition tasks with fewer training iterations compared to classical counterparts.

4.6. Material science and energy

Quantum simulations aid in designing novel materials, such as high-temperature superconductors or efficient photovoltaic cells. For example, IBM's quantum team modeled the electronic structure of lithium-ion battery components to identify pathways for improving energy density—a critical step toward sustainable energy storage.

5. Challenges and future prospects

5.1. Decoherence mitigation

Phonon-induced tunneling fluctuations limit qubit coherence times. Solutions include:

- Hyperbolic metamaterials to suppress phonon density of states.
- Error-corrected algorithms compensating for tunneling errors.

5.2. Sustainable nanoelectronics

Tunnel junctions in neuromorphic chips reduce energy consumption by 100x compared to CMOS. However, rare-material usage (e.g., Pt in STM tips) necessitates recycling strategies.

6. Conclusion

Electron quantum tunneling bridges the quantum and classical worlds, enabling technologies once deemed impossible. As we push the limits of ultrafast control and correlated material design, tunneling will drive innovations in post-Moore's law electronics and topological quantum matter. Interdisciplinary collaboration—spanning condensed matter physics, machine learning, and sustainability science—is essential to fully harness this quantum phenomenon.

Quantum computing represents a transformative leap in computational science, offering unprecedented capabilities to tackle problems that are fundamentally intractable for classical systems. By leveraging quantum superposition, entanglement, and ultrafast control dynamics, this technology has demonstrated potential across cryptography, material science, and optimization. However, the path toward scalable, fault-tolerant quantum computers remains fraught with challenges, including decoherence mitigation, error correction at scale, and the integration of quantum-classical hybrid architectures.

This work underscores the interdependence of theoretical advancements, experimental innovations, and cross-disciplinary collaboration in addressing these barriers. While near-term applications (e.g., quantum-enhanced optimization and molecular simulations) are already emerging, achieving universal quantum supremacy will require breakthroughs in qubit coherence times, high-fidelity gate operations, and novel algorithmic frameworks. Future research should prioritize the development of robust quantum error-correcting codes, scalable fabrication techniques for qubit arrays, and standardized benchmarks for quantifying quantum advantage.

Ultimately, the societal impact of quantum computing hinges not only on technological progress but also on ethical foresight and global cooperation. As the field transitions from laboratory prototypes to real-world deployment, proactive engagement with policymakers and industry

DOI: 10.54254/2753-8818/2025.GL33604

stakeholders will be critical to harnessing its benefits while mitigating risks such as cryptographic vulnerabilities and resource inequity.

References

[1] Esaki, L. (1958). New Phenomenon in Narrow Germanium p-n Junctions. Physical Review.
[2] Binnig, G., Rohrer, H. (1986). Scanning Tunneling Microscopy. IBM Journal of R&D.
[3] Tinkham, M. (2004). Introduction to Superconductivity. Dover Publications.
[4] Goulielmakis, E. et al. (2017). Real-time observation of valence electron motion. Nature.
[5] Zhao, K. et al. (2023). High-fidelity spin qubits via tunneling noise suppression. Nature Nanotechnology.