Proceedings of the 4th International Conference on Applied Physics and Mathematical Modeling
DOI: 10.54254/2753-8818/2026.36902

From Symmetry to Standard Model: Development and Experimental Verification of Yang-Mills Gauge Theory

Yuting Ye

Department of Physics, University of Michigan, Ann Arbor, USA
yutingye@umich.edu

Abstract. This article examines how gauge symmetry developed from a principle of invariance into the organizing framework of the Standard Model. It focuses on the transition from global to local symmetry, the Yang-Mills generalization to non-Abelian groups, and the application of gauge theory to quantum chromodynamics and electroweak unification. Using conceptual analysis, historical reconstruction, and comparison with landmark experimental evidence, the study explains why local transformations require gauge connections, how non-commuting fields generate self-interactions, and how the Higgs vacuum gives mass to weak gauge bosons. The analysis then evaluates neutral-current scattering, the discoveries of the W and Z bosons, and the observation of the Higgs boson as tests of the framework. These results show that Yang-Mills theory successfully anticipated essential particles and interactions while remaining incomplete in areas such as neutrino mass, dark matter, gravity, and the strong-CP problem. The paper concludes that precision Higgs, QCD, neutrino, and electric-dipole measurements offer promising routes toward physics beyond the Standard Model.

Keywords: Gauge Symmetry, Yang–Mills Theory, Standard Model, Electroweak Unification, Higgs Mechanism

1. Introduction

Modern particle theory often begins by asking which transformations leave an action unchanged, rather than by postulating a force law directly. Its basic objects are therefore the action, the Lagrangian, and the transformations that preserve them. Noether connected continuous symmetries with conservation laws. Gauge symmetry adds a stronger requirement: allowing the transformation to vary over spacetime restricts the form of an interaction. In electromagnetism, making the $U(1)$ phase of a fermion field position dependent spoils ordinary differentiation; covariance is restored by introducing a gauge connection. Yang and Mills showed in 1954 that the same logic could be applied to a non-Abelian internal group [1]. Because the generators do not commute, the resulting gauge fields couple to one another, unlike the photon in Abelian electrodynamics. This feature eventually made Yang-Mills theory the mathematical basis of the Standard Model, whose gauge symmetry is $SU(3)_C \times SU(2)_L \times U(1)_Y$. QCD supplies its strong-interaction sector, whereas the electroweak construction places electromagnetic and weak processes in one gauge description [2,

© 2026 The Authors. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (https://creativecommons.org/licenses/by/4.0/).

DOI: 10.54254/2753-8818/2026.36902

3]. A nonzero Higgs vacuum then gives mass to selected gauge bosons without abandoning the gauge construction. After developing the global-to-local argument, this paper applies it to QCD and electroweak unification and then compares the resulting model with experiment, highlighting the discovery of weak neutral currents, $W^{\pm}$, Z bosons, and the Higgs boson, before considering precision Higgs studies, QCD measurements, and the strong-CP puzzle as possible windows on a more complete theory.

2. From symmetry to gauge theory

A field theory can be organized by the symmetries of its action instead of by a catalogue of pairwise forces. The action formulation is valuable because it puts dynamics and invariance in a single mathematical setting. Noether's theorem makes this connection precise: invariance under continuous changes produces conserved currents. In particular, time- and space-translation symmetry account for conservation of energy and momentum. The important distinction is whether the same transformation is used everywhere. A global symmetry uses one common parameter, whereas a gauge symmetry permits a different choice at every spacetime point and consequently demands additional mathematical structure. Action is defined as $S=\int d^4x\,\mathscr{L}$, S is the action, $\mathscr{L}$ is the Lagrangian, d^4x is integrating over the four-dimensional space-time. Setting the first variation of this action to zero produces the equations obeyed by the fields.

The Lagrangian for free fermion field is $\mathscr{L}_0=\bar{\psi}\,(i\gamma^\mu\partial_\mu-m)\psi$. ψ is the fermion field, $\bar{\psi}=\psi^\dagger\gamma^0$ is the Dirac conjugate, m is the particle mass, γ^μ are the gamma matrices. Consider the global phase transformation $\psi(x)\to e^{iq\alpha}\psi(x)$. At the same time, the Dirac conjugate field transformation $\bar{\psi}(x)\to\bar{\psi}(x)e^{-iq\alpha}$. Here, α is a constant independent of spacetime position. Since the phase factors cancel each other out in the Lagrangian, the free Dirac Lagrangian remains invariant under this transformation. According to Noether's theorem, there is a correspondence between continuous global $U(1)$ symmetry and a conserved current $j^\mu=\bar{\psi}\gamma^\mu\psi$ and $\partial_\mu j^\mu=0$. Multiplying the fermion field everywhere by one common phase does not alter measurable probabilities. The global $U(1)$ operation thus represents freedom in the choice of overall phase, and Noether's theorem associates that continuous freedom with charge conservation. For a uniform rotation, the parameter is constant and differentiation introduces no extra contribution, so no compensating vector field is needed. That conclusion fails once the phase is permitted to vary with position.

If the constant parameter α is replaced by a function $\alpha(x)$ that depends on the spacetime position, the local phase transformation must be rewritten as $\psi(x)\to e^{iq\alpha(x)}\psi(x)$. Since the phase now depends on spacetime coordinates, the transformation of ordinary derivatives gives rise to extra terms $\partial_\mu\psi\to e^{iq\alpha(x)}\left[\partial_\mu+iq\,\partial_\mu\alpha(x)\right]\psi$. And this extra term $iq\,\partial_\mu\alpha(x)$ break the local invariance of the original Lagrangian. Restoring covariance requires an additional vector field whose transformation compensates for the derivative of the local phase; ordinary differentiation is therefore replaced by covariant differentiation. A_μ $D_\mu=\partial_\mu+iqA_\mu$. Then, the transformation of the gauge field needs to be rewritten as $A_\mu\to A_\mu-\partial_\mu\alpha$ make $D_\mu\psi\to e^{iq\alpha(x)}D_\mu\psi$. The Lagrangian for the interaction between fermions and the electromagnetic field can be written as $\mathscr{L}=\bar{\psi}\,(i\gamma^\mu D_\mu-m)\psi-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ and $F_{\mu\nu}=\partial_\mu A_\nu-\partial_\nu A_\mu$. A local phase convention assigns a

DOI: 10.54254/2753-8818/2026.36902

separate reference at neighboring points. An ordinary derivative compares the fields at those points without accounting for the change of convention, which is why it fails to transform covariantly. A connection is needed to define that comparison consistently. The gauge potential A_μ supplies this connection, and the covariant derivative corrects the field's ordinary variation for the changing local phase reference. On this reading, the electromagnetic potential is not only a convenient route to electric and magnetic fields; it is the connection that makes comparisons between local phase choices meaningful.

3. Yang–Mills theory and non-Abelian gauge fields

The above describes electromagnetic $U(1)$ gauge theory. The central step taken by Yang and Mills was to show that local gauge invariance need not be restricted to the Abelian $U(1)$ symmetry of electromagnetism [1]. For a general non-Abelian gauge group G, the local gauge transformation is written as $\psi(x) \to U(x)\psi(x)$ and the local group element is $U(x) = exp\left[ig\alpha^a(x)T^a\right].T^a$ are the generators of the Lie group, $\alpha^a(x)$ are the local transformation parameters, g is the gauge coupling constant, the index a labels the different group generators and $\left[T^a, T^b\right] = if^{abc}T^c$. Here, f^{abc} are the structure constants of the group. Because each independent generator corresponds to a direction in the internal symmetry space, the local theory requires a corresponding set of gauge-field components. A_μ^a The covariant derivative is written as $D_\mu = \partial_\mu - igA_\mu^aT^a$. Define $A_\mu = A_\mu^aT^a$ and make $D_\mu = \partial_\mu - igA_\mu$, then the Yang-Mills field strength can be defined via the commutator of the covariant derivative $[D_\mu, D_\nu] = -igF_{\mu\nu}$ and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ig\left[A_\mu, A_\nu\right]$. This can also be written as $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^bA_\nu^c$, $gf^{abc}A_\mu^bA_\nu^c$. The final term is present because non-Abelian generators fail to commute. It makes the field strength nonlinear and allows gauge quanta to interact directly with other gauge quanta. The Lagrangian of the Yang-Mills gauge field is $\mathscr{L}_{YM} = -\frac{1}{4}F_{\mu\nu}^aF^{a\mu\nu}$. This is not equivalent to placing several independent electromagnetic fields side by side. The mediator carries the non-Abelian charge and therefore enters its own dynamics, producing self-couplings without a QED counterpart. Such couplings are indispensable for the behavior of both the weak gauge fields and gluons [1].

The gauge group of the Standard Model is $G_{SM} = SU(3)_C \times SU(2)_L \times U(1)_Y$. While $SU(3)_C \longrightarrow$ strong interaction and $SU(2)_L \times U(1)_Y \longrightarrow$ electroweak interaction. Glashow, Salam, and Weinberg developed a common gauge description for weak and electromagnetic processes. The Brout-Englert-Higgs construction then explained massive weak carriers through their interaction with a nonzero vacuum field rather than through explicit gauge-boson mass terms. Quantum Chromodynamics (QCD), with $SU(3)$ as its gauge group, is a type of Yang–Mills theory. The discovery of asymptotic freedom around 1973 revealed that the effective interaction between quarks weakens extremely short distances and high energies [2, 3], At sufficiently high momentum, this weakening makes perturbative calculations reliable and identifies QCD as the Standard Model theory of strong processes.

4. Applications of Yang-Mills theory in the standard model

Among the Standard Model sectors, QCD displays the distinctive consequences of non-Abelian gauge dynamics most directly. QCD describes the color degrees of freedom of quarks in terms of a local $SU(3)_C$ gauge symmetry. Quarks have three colors $q = (q_r, q_g, q_b)^T$. A gluon carries color

DOI: 10.54254/2753-8818/2026.36902

charge, whereas a photon carries no electric charge. Gluons can therefore couple to one another, as required by the noncommuting generators of $SU(3)$ group. Gross, Wilczek, and Politzer showed that the QCD interaction becomes progressively weaker when probed at higher momentum scales, the behavior now known as asymptotic freedom [2, 3]. The development of QCD is closely linked to particle physics research in the 1970s. Early studies revealed a rapidly growing number of hadrons, making it difficult to explain the underlying patterns through simple particle classification alone. A major step toward the modern description of strong interactions came in 1964, when Gell-Mann and Zweig independently introduced quarks as underlying constituents of hadrons. However, the quark model was initially merely a classification scheme and did not explain the nature of the interactions between quarks. With the development of Yang-Mills theory, physicists realized that the strong interaction could be naturally described as an $SU(3)_C$ color gauge theory. Gross, Wilczek, and Politzer supplied the decisive theoretical result: QCD is asymptotically free, so quarks interact weaklier at short distances. This behavior connected the color gauge theory to high-energy scattering data.

Electromagnetic and weak processes appear unrelated at ordinary energies: one has macroscopic range, whereas the other is confined to subatomic distances. Electroweak theory attributes this contrast to the broken, low-energy phase of a shared symmetry. Glashow, Weinberg, and Salam formulated the electroweak unification theory, combining $SU(2)_L$ and $U(1)_Y$. When electroweak symmetry is preserved $W_\mu^1, W_\mu^2, W_\mu^3, B_\mu$ are all massless gauge fields. After electroweak symmetry breaking, the electromagnetic gauge field γ remains associated with a massless photon, whereas the weak fields form the massive $W^\pm$ and states Z. The equations remain gauge invariant even though the vacuum preserves only the electromagnetic subgroup. In this way the theory accommodates massive weak carriers without the inconsistency created by inserting their masses by hand.

The development of the electroweak unification theory stemmed from a long-standing issue: although electromagnetic and weak interactions appear completely different in low-energy experiments, both involve interactions between leptons and gauge fields. In the 1950s, Glashow first proposed describing the weak interaction using a gauge theory with higher symmetry, subsequently, Weinberg and Salam independently developed electroweak models incorporating the Higgs mechanism, which allowed the gauge bosons of the weak interaction to acquire mass while preserving the theory's quantum consistency. The theory's most significant contribution was demonstrating that photons and W, Z bosons do not originate from entirely different sources but rather emerge from a unified electroweak gauge structure at higher energy scales. Spontaneous breaking thus generates weak-boson masses from the vacuum while leaving explicit mass terms out of the original gauge Lagrangian. In the 1960s, several physicists including Brout, Englert, Higgs, and Guralnik, Hagen, and Kibble independently proposed similar ideas. They discovered that the mass of gauge bosons did not need to be acquired through the direct addition of a mass term but could instead arise from a change in the vacuum structure. This mechanism subsequently became crucial to the viability of electroweak theory. In the 1970s, the proof of the renormalizability of electroweak theory further bolstered the theory's credibility.

The Standard Model assigns half-integer-spin fields to matter and integer-spin fields to force mediation or symmetry breaking. Six quark flavors and six leptons form the fermion sector; the boson sector consists of gluons, the photon, $W^\pm, Z$, together with the Higgs boson. Matter fields couple through the gauge bosons, whose identities and allowed interactions follow from their group representations. The classification is therefore structural, not merely a list assembled from observations.

DOI: 10.54254/2753-8818/2026.36902

Some of the model's strongest evidence comes from phenomena predicted before they were measured. In 1973, The Gargamelle experiment at CERN discovered weak neutral currents. In 1983, CERN's UA1 and UA2 experiments discovered $W^{\pm}, Z$. In 2012, the LHC's ATLAS and CMS experiments discovered the Higgs boson [4, 5]. Together, these observations show that the gauge construction has empirical content beyond its mathematical economy. Its experimental record does not make the Standard Model complete. Gravity lies outside its gauge structure, and its known particles do not account for the dark matter and dark energy inferred from cosmology. Furthermore, certain parameters of the Standard Model, such as the mass hierarchies of different fermions and the strong CP problem still lack a deeper theoretical explanation. The theory is therefore both the benchmark for particle physics and the baseline against which evidence for a deeper framework must be judged. Many current high-energy physics experiments continue to seek new phenomena beyond the scope of existing gauge theories, such as supersymmetry, extended gauge structures, and dark matter candidates.

5. Experimental verification and theoretical success of the standard model

In 1973, Gargamelle recorded neutrino-scattering events in which no electric charge was transferred, providing the first evidence for a neutral weak process. As shown in Figure 1, the bubble-chamber photograph records the characteristic particle tracks used to identify such interactions [6, 7].

Traditional weak interactions are charged-current processes, $\nu + N \rightarrow \nu + N$. In the case of weak neutral currents, the neutrino remains a neutrino after the collision and carries no electric charge. This event class was a nontrivial test because the electroweak model specifically required a neutral current in addition to the familiar charged current. The weak interaction involves not only the $W^{\pm}$ bosons but also the theoretically predicted neutral gauge boson.




Figure 1. Neutrino tracks in the CERN Gargamelle bubble chamber [7]

The same framework also required heavy charged and neutral weak mediators, identified as the $W^{\pm}$ and bosons Z . Producing these heavy mediators required collider energies far above those associated with electrons or protons at rest. UA1 and UA2 obtained direct evidence at CERN in

DOI: 10.54254/2753-8818/2026.36902

1983 for the $W^{\pm}, Z$. Their measured masses agreed closely with the electroweak predictions. $m_W \approx 80.4\,\mathrm{GeV}$ $m_Z \approx 91.2\,\mathrm{GeV}$.

Although the electroweak theory had successfully predicted the W and Z bosons before they were found, the scalar sector remained unverified. In the mechanism, the vacuum has a nonzero scalar-field value, and an excitation about that value appears experimentally as the Higgs boson. Independent ATLAS and CMS data sets reported in 2012 showed a new boson near $m_H \approx 125\,GeV$.

Both experiments observed a signal with a significance exceeding 5σ. Particle-physics experiments conventionally require a five-standard-deviation significance before claiming a discovery [4, 5].




Figure 2. Mass distributions of Higgs boson candidate events in the ATLAS and CMS experiments [4, 5]

As shown in Figure 2, the concentration of Higgs-boson candidate events near 125 GeV exceeded the predicted background. This localized excess was an essential component of the evidence that the observed state was a new boson.

6. Precision tests and new physics beyond the standard model

After the Higgs discovery, the main task changed from filling a missing entry in the particle spectrum to testing the model's parameters more stringently [6, 7]. Collider programs now pursue two complementary strategies: direct searches for new states and indirect searches for statistically stable departures in well-measured quantities. It includes precision electroweak measurements, such as W boson mass, Z boson properties, and the weak mixing angle. QCD tests and experimental verification of the strong interaction: jet production, the running of the strong coupling constant, and gluon interactions [8, 9].

Although experiments have repeatedly confirmed the Standard Model, contemporary high-energy physics is increasingly concerned with small, reproducible discrepancies rather than with completing its original particle inventory. Many observables can now be predicted with enough accuracy that even a modest but persistent disagreement could point to dynamics outside the

DOI: 10.54254/2753-8818/2026.36902

established theory. The Higgs sector is especially informative because it is still known less precisely than the gauge sector. Since 2012, ATLAS and CMS have moved from establishing that the particle exists to measuring how often it is produced, how it decays, and how strongly it couples to other particles. A consistent departure in any of these quantities could signal extra scalar degrees of freedom or a modified gauge structure [10, 11]. Neutrinos provide a separate indication that the minimal model is incomplete. Oscillation measurements show that neutrino flavor changes during propagation, which requires nonzero neutrino masses and therefore contradicts the massless-neutrino assumption of the original formulation [12, 13]. Determining how those masses arise is now one of the principal experimental routes beyond the Standard Model.




Figure 3. Results from the ATLAS experiment on the coupling strengths of the Higgs boson to various particles [6]

Figure 3 shows the measured Higgs-boson coupling strengths. Within present experimental errors, the measured couplings remain compatible with Standard Model expectations. Reducing those errors may therefore be more revealing than simply repeating the original discovery tests.

Future measurements will also target the Higgs field's self-interaction, not only its couplings to fermions and gauge bosons [10, 11]. In the Standard Model, the form of the scalar potential determines the Higgs self-coupling, so these observable tests the symmetry-breaking sector directly. Its value is still poorly constrained. A measured self-coupling inconsistent with the prediction would suggest that electroweak symmetry breaking involves more than the single scalar field assumed by the minimal theory. The Higgs is therefore no longer merely a discovered particle; it is an experimental probe of the mechanism that selects the electroweak vacuum. In gauge-theory terms, precision Higgs measurements test whether the broken phase behaves as the Standard Model says it should.

Beyond the precision experimental testing of the Standard Model, there remain unresolved issues within QCD. A particularly puzzling unresolved feature of QCD concerns its apparent preservation of CP symmetry. According to Yang-Mills theory, the $SU(3)_C$ gauge field allows for an additional topological term that leads to CP symmetry breaking in the strong interaction [14, 15]. CP combines charge conjugation, which interchanges particles and antiparticles, with spatial inversion. Neutral K and B mesons demonstrate that weak processes can violate this symmetry. QCD also permits CP

DOI: 10.54254/2753-8818/2026.36902

violation mathematically, yet no comparably large strong-interaction effect has been observed. The puzzle is the gap between that allowed QCD term and the exceptionally small experimental bound on its physical consequences. Searches for a neutron electric dipole moment constrain the strong CP parameter to lie extremely close to zero [14]. Researchers also proposed a new global symmetry that dynamically relaxes the offending parameter; breaking that symmetry produces an axion [15]. An axion would both solve the strong-CP problem and supply a concrete target for physics beyond the Standard Model. Ongoing searches therefore connect a formal feature of Yang-Mills theory with present experimental programs.

7. Conclusion

The move from Abelian invariance to the Yang-Mills construction turned symmetry from a classification tool into a rule for building interactions. The Standard Model uses the non-Abelian product group $SU(3)_C \times SU(2)_L \times U(1)_Y$ to describe strong and electroweak processes, with the Higgs vacuum accounting for massive weak carriers. Neutral-current events and the subsequent discoveries of $W^{\pm}, Z$ and the Higgs particle supplied distinct experimental checks of this construction. The agreement is impressive but incomplete. More accurate measurements of Higgs couplings, Higgs self-interaction, and QCD observables may expose small failures of the current predictions, while the strong-CP problem already points to missing theoretical structure. Yang-Mills theory consequently has a dual role: it explains the interactions that experiments have confirmed and provides the framework in which many proposed extensions are expressed.

Looking ahead, higher-luminosity colliders, improved lattice-QCD calculations, precision neutrino experiments, and sensitive searches for electric dipole moments may reveal where the Standard Model fails. Progress will likely come from combining increasingly accurate gauge-theory predictions with complementary measurements, guiding extensions involving new symmetries, scalar fields, axions, or dark-sector particles and dynamics.

References

[1] Chevyrev, I. (2022). Stochastic quantization of Yang–Mills. *Journal of Mathematical Physics, 63*(9), 091101.
[2] Gross, D. J., & Wilczek, F. (1973). Ultraviolet behavior of non-Abelian gauge theories. *Physical Review Letters, 30*(26), 1343–1346.
[3] Politzer, H. D. (1973). Reliable perturbative results for strong interactions? *Physical Review Letters, 30*(26), 1346–1349.
[4] ATLAS Collaboration. (2012). Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Physics Letters B, 716*(1), 1–29.
[5] CMS Collaboration. (2012). Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Physics Letters B, 716*(1), 30–61.
[6] ATLAS Collaboration. (2022). A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery. *Nature, 607*, 52–59.
[7] CMS Collaboration. (2022). A portrait of the Higgs boson by the CMS experiment ten years after the discovery. *Nature, 607*, 60–68.
[8] Potrebko, A. (2024). Review of the measurements of the strong coupling constant in CMS at 13 TeV (CMS-CR-2024-092) [Conference report]. CERN. https://cds.cern.ch/record/2898147.
[9] Connor, P. L. S. (2024). Determination of the strong coupling from jet measurements at CMS. *Proceedings of Science, ICHEP2024*, 501.
[10] Dawson, S., et al. (2022). Report of the topical group on Higgs physics for Snowmass 2021: The case for precision Higgs physics. arXiv:2209.07510.
[11] Jakobs, K., & Zanderighi, G. (2024). The profile of the Higgs boson: Status and prospects. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences, 382*(2266), 20230087.

DOI: 10.54254/2753-8818/2026.36902

[12] Granelli, A. (2024). Standard and non-standard aspects of neutrino physics. *Universe, 10*(4), 164.
[13] Bhupal Dev, P. S. (2026). Neutrino theory overview. In *Proceedings of the XXVI DAE-BRNS High Energy Physics Symposium 2024* (pp. 13–19). Springer.
[14] Roussy, T. S., Caldwell, L., Wright, T., Cairncross, W. B., Shagam, Y., Ng, K. B., Schlossberger, N., Park, S. Y., Wang, A., Ye, J., & Cornell, E. A. (2023). An improved bound on the electron's electric dipole moment. *Science, 381*(6653), 46–50.
[15] Sannino, F. (2026). Strong CP and the QCD axion lecture notes via effective field theory. *The European Physical Journal C, 86*, 465.